




THE CONSERVATION STATUS OF PICABO MILKVETCH (*ASTRAGALUS ONICIFORMIS* BARNEBY)

159-13

by
Robert K. Moseley
and Steve J. Popovich

PLANT GENERALLY PROSTRATE,
2-6 INCHES TALL, UP TO 2
FEET IN DIAMETER



PICABO MILKVETCH
(*ASTRAGALUS ONICIFORMIS*)

QL
84.2
.L352
no. 95-9



#35336413

ID 8806A232

QL
84.2
.L352
no. 95-9

THE CONSERVATION STATUS OF
PICABO MILKVETCH (*ASTRAGALUS ONICIFORMIS* BARNEBY)

by

Robert K. Moseley
Conservation Data Center
Idaho Department of Fish and Game
600 S. Walnut, P.O. Box 25
Boise, ID 83712

and

Steve J. Popovich
Shoshone District
Bureau of Land Management
400 West F St., P.O. Box 2-B
Shoshone, ID 83352

January 1995



ABSTRACT

Astragalus onitiformis Barneby (Picabo milkvetch) is a narrowly-distributed endemic to the north-central portion of the eastern Snake River Plain. Habitat for nearly the entire known distribution of this species is managed by the BLM, mostly the Shoshone District, but also the Idaho Falls District. Due to its rarity, it is a federal category 3c candidate and a BLM sensitive species. Intensive searches, beginning in 1981, have resolved the distribution, abundance, and conservation status of the species. Thirty-six occurrences, consisting of numerous subpopulations, occur within a main range of about 42 x 13 miles, with several disjunct populations lying outside the core of its distribution. Short- and long-term threats have been identified. Any conservation strategy developed to assure the long-term persistence of Picabo milkvetch as a viable evolutionary unit must take into account the decline of the sagebrush-steppe ecosystem on the Snake River Plain.



TABLE OF CONTENTS

ABSTRACT	i
TABLE OF CONTENTS	ii
LIST OF APPENDICES	ii
INTRODUCTION	1
RESULTS	1
<i>ASTRAGALUS ONICIFORMIS</i>	2
Taxonomy	2
Legal or Other Formal Status	2
Description	3
Distribution	5
Habitat	7
Population Biology	10
Assessment and Management Recommendations	14
REFERENCES	17

LIST OF APPENDICES

- Appendix 1 ... Line drawings of *Astragalus oniciformis*.
- Appendix 2 ... Maps of the distribution of *Astragalus oniciformis*.
- Appendix 3 ... Occurrence records from the Conservation Data Center for *Astragalus oniciformis*.



INTRODUCTION

Picabo milkvetch (*Astragalus oniciformis* Barneby) is a narrow endemic, restricted to sandy soils in the north-central portion of the eastern Snake River Plain, Idaho. It has been recognized to be of conservation concern since the early 1970's (Johnson and Steele 1974), and as BLM sensitive species for a number of years (e.g., Conservation Data Center 1994). It was once considered a candidate for listing under the Endangered Species Act, first as a category 1 species (U.S. Fish and Wildlife Service 1980), and later as a category 2 candidate (U.S. Fish and Wildlife Service 1983; 1985). It was later downgraded to a category 3c candidate (U.S. Fish and Wildlife Service 1990). Category 3c is used for those candidate species that are later found to be more common or under less threat than previously thought. This reconsideration by the U.S. Fish and Wildlife Service was largely a result of recommendations of Packard and Smithman (1984) who performed the first systematic field inventory for Picabo milkvetch in 1984. In the intervening decade since the last field survey, a large portion of Picabo milkvetch habitat has been altered by wildfire and range improvement seedings and the current status of its distribution and abundance are unknown.

Most Picabo milkvetch habitat occurs on public lands managed by the Shoshone District BLM. Because land management activities could have a major impact on the long-term viability of the species, the BLM felt a current status inventory should be undertaken. Aside from mapping populations they discovered and collecting numerous voucher specimens, Harrison (1981) and Packard and Smithman (1984) collected little population and habitat information. To rectify this paucity of information on the current conservation status of Picabo milkvetch, the Shoshone District BLM and the Idaho Department of Fish and Game's Conservation Data Center (CDC) entered into a cooperative project to conduct field inventories in 1994. The primary objectives of this investigation are as follows:

- 1) Survey and delineate known populations of Picabo milkvetch, and search for additional populations.
- 2) Characterize habitat conditions for the populations.
- 3) Assess population data and threats to the species and make management recommendations to the Shoshone District BLM based on these assessments.

RESULTS

During May and June 1994, Moseley conducted a field survey of nearly all suitable habitat west of the Craters of the Moon Lava Field and Wapi Lava Field. Popovich conducted a survey for the species east of the Craters of the Moon Lava Field in June 1994 (Popovich 1994a), filling in much needed distribution data. Following is the status of our knowledge of the distribution, abundance, and conservation status of Picabo milkvetch, including information on taxonomy, habitat, distribution, conservation status, and management and conservation recommendations. Line drawings of Picabo milkvetch, distribution maps, and occurrence records are appended to the end of the report.

Astragalus oniciformis Barneby

TAXONOMY

Bibliographic citation: Barneby, R.C. 1957. *Pugillus Astragalorum* XX: Notes on *A. mulfordae* [sic] and some close relatives. Leaflets in *Western Botany* 8:120-125.

Type specimen: Idaho, Blaine County, elevation 4750 ft, east of Picabo, 21 June 1947.

Pertinent synonym(s): None.

Common name: Picabo milkvetch.

Size of genus: A vast genus of perhaps 2000 species, most highly developed in arid continental, desert, and Mediterranean climates, circumboreal in dispersal, most numerous in central Asia, Iran and Turkey, in western North America, and in the Andes of South America (Barneby 1964).

Family name: Fabaceae; Leguminosae

Common name for family: Pea.

History of knowledge of taxon in Idaho: Although previously collected by Nelson and Macbride, and possibly Ray Davis, Rupert Barneby collected this species near the town of Picabo in 1947. He did not describe it as a new species, however, until 1957 (Barneby 1957). For over two decades, Picabo milkvetch was known only from this locality (Barneby 1964; Steele 1975; Packard 1981), although other sources say that it was "plentiful" in the foothills of the Sawtooth Mountains, Blaine County (Hitchcock 1961; Hitchcock and Cronquist 1973). Early rare plant inventories of the Shoshone District failed to locate any populations (Eidemiller 1977a; 1977b). It was not until Harrison's (1981) survey for the BLM in 1981, that any new populations were located. Packard and Smithman (1984) were the first to conduct a systematic inventory for the species and, with a few slight modifications by Ann DeBolt and Steve Popovich, the distribution for Picabo milkvetch that they delimited stood until 1994.

Alternative taxonomic treatments: None.

LEGAL OR OTHER FORMAL STATUS

National:

U.S. Fish and Wildlife Service: Picabo milkvetch is a category 3c candidate for listing under the Endangered Species Act (U.S. Fish and Wildlife Service 1990). Category 3c includes those taxa that have proven to be more abundant or widespread than previously believed and/or those that are not subject to any identifiable threat. If further research or changes in habitat indicate a significant decline in these taxa, they may be reevaluated for possible inclusion in categories 1 or 2 (U.S. Fish and Wildlife Service 1993).

Bureau of Land Management: Picabo milkvetch is currently an Idaho BLM Sensitive Species (Conservation Data center 1994).

Other current formal status recommendations: Because it is rare and uncommon, but not currently imperiled, it is given a global (G) conservation rank of 3 (on a scale of 1 to 5) by the Association for Biodiversity Information (the International Association of Natural Heritage Programs and Conservation Data Centres) (Conservation Data Center 1994).

State: (Picabo milkvetch is endemic to Idaho.)

Idaho:

Idaho Native Plant Society: Picabo milkvetch is a Monitor species on the Idaho Native Plant Society list of the state's rare flora. The Monitor list includes species that are common within a limited range in Idaho, and/or are uncommon, but have no identifiable threats (Idaho Native Plant Society 1994).

Conservation Data Center: Because Picabo milkvetch is endemic to Idaho, the state (S) conservation rank assigned by the Conservation Data Center (the Idaho node of the Association for Biodiversity Information) equals the global (G) rank of 3, discussed above (Conservation Data Center 1994).

Review of past status: In her review of this taxon for the Rare and Endangered Plants Technical Committee of the Idaho Natural Areas Council, Packard (1981) recommended a federal status of Endangered, admitting, however, that further study was needed. Harrison (1981) also recommended Endangered status.

DESCRIPTION

General nontechnical description: Picabo milkvetch is a wiry, prostrate plant with numerous stems emanating from a loosely branching caudex at ground level. The caudex surmounts a rather long, slender taproot. The stems are from 1 to 2.5 dm long, sparsely leafy, and only the lowest leaf axils without an inflorescence. Leaves have a short petiole, or the upper ones nearly without a petiole, a stiff rachis, and 17 to 25 small, scattered, oval leaflets. The herbage is covered with short, white hairs. A raceme of 6-12, cream-yellow, pea-like flowers comprises the inflorescence. The small fruit is pendulous, attached by a short stipe, light green-papery, and more or less triangular in cross section.

Technical description: Slender but wiry, diffuse, perennial, with a taproot and at length loosely forking, suffruticose caudex, loosely strigulose or subvillosulous with incumbent or ascending, nearly straight or sinuous hairs up to 0.35-0.5 (0.6) mm long, the herbage greenish-cinereous, the leaflets glabrous or medially glabrescent above; stems several or numerous, decumbent or prostrate, (0.5) 1-2.5 dm long, commonly spurred at the lowest nodes and thence floriferous upward from all succeeding axils; stipules papery-membranous or early becoming so, 1.5-3.5 (4) mm long, triangular or triangular-acuminate, decurrent around half, or the lowest around the whole stem's circumference but not connate; leaves 2.5-7.5 cm long, shortly petioled or the upper ones subsessile, with stiff but

slender rachis and (13) 17-25 (27) rather distant, scattered, narrowly to broadly elliptic, oval or oblong, obtuse to retuse, mostly folded leaflets 1-6.5 mm long; *peduncles* 0.5-2.5 cm long, much shorter than the leaves; *racemes* very loosely (4) 6-12-flowered, the flowers early nodding, the axis becoming 1-7 cm long in fruit, usually produced beyond the last flower as a subulate appendage; *bracts* papery-membranous, ovate or lanceolate, 0.5-1.8 mm long; *pedicels* filiform, very early arched outward or ultimately recurved, at anthesis 1-1.4 mm, in fruit 1-1.6 mm long, tardily disjointed with the fruit; *bracteoles* 0; *calyx* 2.9-3.8 mm long, strigulose-villosulous like the herbage with white or some dark hairs, the subsymmetric disc 0.6 mm deep, the campanulate tube 2.1-2.3 mm long, 1.7-2 mm in diameter, the subulate teeth 0.8-1.5 mm long, the whole becoming papery, marcescent unruptured; *petals* ochroleucous, the banner veined with dull brownish-lilac; *banner* abruptly recurved through 90° or sometimes further, broadly ovate cuneate, 5.3-6.8 mm long, 4-5 mm wide; *wings* (0.7 mm longer to 0.7 mm shorter than the banner) 5.7-6.8 mm long, the claws 2-2.5 mm, the oblong-obovate, obtuse blades 4-4.4 mm long, 1.9-2.9 mm wide, both incurved but the left one further and more abruptly than the right; *keel* 4-5 mm long, the claws 1.8-2.3 mm, the half-circular blades 2.4-3 mm long, 1.7-1.9 mm wide, abruptly incurved through 110-130° to the deltoid, often obscurely correct apex; *anthers* 0.35-0.5 mm long; *pod* pendulous, stipitate, the stipe (1.5) 2-4 mm long, the body lance- or oblong-ellipsoid, usually a trifle incurved, 7-12 mm long, (2) 2.5-3.5 (3.8) mm in diameter, cuneate at both ends or truncately contracted at the base into the stipe, cuspidate at apex, triquetrously compressed with nearly flat lateral and openly sulcate dorsal faces, keeled ventrally by the prominent suture, the thin, green but rather densely strigulose valves becoming papery, stramineous, delicately cross-reticulate, inflexed as a not quite complete septum 0.7-1.2 mm wide; *dehiscence* apical and through the ventral suture; *ovules* 6-12; *seeds* brown or greenish-brown, sometimes purple-speckled, smooth but dull, 1.5-1.8 mm long (Barneby 1964).

Local field characters: The elegantly pretty Picabo milkvetch is easily picked out from among other species on the eastern Snake River Plain with small, whitish to yellowish flowers. It is recognized by its combination of wiry, prostrate stems, and its pod that is pendulous, stipitate, and almost exactly triquetrous, the three faces being of almost equal width. The pod is sometimes a trifle turgid, and the papery valves become straw-colored and shiny in ripening. Neither of the two other *Astragali* known to be sympatric with Picabo milkvetch, *Astragalus purshii* and *A. lentiginosus*, share these characteristics. Another microphyllous milkvetch, also largely endemic to the Shoshone District BLM, that may cause some confusion is *Astragalus atratus* var. *inseptus* (Eidemiller 1977a; 1977b). The main characteristic that differentiates it from Picabo milkvetch is the presence of a pod that is usually purplish or mottled when mature and is not conspicuously triquetrous. Other morphological and ecological differences also exist (Smithman 1989).

In many respects, Picabo milkvetch resembles *Astragalus mulfordiae*, with which it is not sympatric, being separated by at least 90 miles [Isely (1986) erroneously states that the two have contiguous distributions]. In fact, Barneby (1957; 1964) referred to this likeness in naming Picabo milkvetch. The specific epithet, *oniciformis*, notes the plant's resemblance to *Astragalus mulfordiae*, which was referred by Rydberg to the genus *Onix*. Care must be taken to distinguish specimens of the two species in the herbarium; there has been confusion in the past. The following key, adapted from Barneby (1964:459), can be used to distinguish the two species:

1. Stipules mostly free; vesture of sinuous or incurved hairs; petals ochroleucous, the banner veined with brownish-purple; leaflets oblong-elliptic, all jointed; body of the pod 6-12 mm long; ovules 6-12; seeds 1.5-1.8 mm long; Blaine, Lincoln, and Minidoka counties, between 3700 and 5200 feet in elevation *Astragalus oniciformis*

1. Stipules connate; vesture of appressed, straight hairs; petals whitish; leaflets mostly linear, the terminal one of some upper leaves nearly always continuous with the rachis; body of the pod (9) 10-16 mm long; ovules (11) 12-16; seeds 2-2.6 mm long; Ada, Owyhee, Payette, and Washington counties, Idaho, and adjacent Malheur County, Oregon; between 2100 and 3200 feet in elevation *Astragalus mulfordiae*

Photos and line drawings: Reproductions of a line drawing of Picabo milkvetch by Jeanne Janish appears in Hitchcock (1961), Hitchcock and Cronquist (1973), Barneby (1989), and Appendix 1. Photographs of its habit and habitat are in the slide collection of the Conservation Data Center.

DISTRIBUTION

Global/Idaho distribution: Picabo milkvetch is endemic to the north-central portion of the eastern Snake River Plain (*sensu* Malde 1991) in Blaine, Lincoln, and Minidoka counties, Idaho (Appendix 2). Although first collected at Picabo, reflected by the common name, that population represents the extreme northwestern corner of its distribution. Using this population as their reference, Barneby (1964; 1989), Hitchcock (1961), and other sources described its range as occurring in the "foothills of the Pioneer Mountains." This has been found to be false; Picabo milkvetch is truly a species of the Snake River Plain.

The CDC data base contains 36 occurrence records for this species. An occurrence record is an information management convenience to track data regarding a population or several subpopulations in a localized area and is identified with the three-digit code (e.g., 012). The general distribution of Picabo milkvetch is mapped in Appendix 2. We also mapped the distribution on two sets of 1:100,000-scale Idaho Transportation Department maps, on file at the CDC and Shoshone District BLM office. Photocopies of these maps have been included in Appendix 2. In addition, the distribution is also mapped on 1:24,000-scale USGS quadrangles on file at the CDC. Picabo milkvetch populations on the latter two maps are referenced by CDC occurrence number. A general description of Picabo milkvetch distribution, defined by its distributional limits, is outlined below:

Northern Limit: The northwestern section of its northern limit follows the base of the foothills of the Pioneer Mountains and Picabo Hills. The foothills are largely underlain by Eocene Challis Volcanics (Rember and Bennett 1979a; Worl et al. 1991) and the soils derived by this bedrock have a clay texture, a habitat unsuitable for Picabo milkvetch (see Habitat section). Apparently little to no sandy loess was deposited in the upper Silver Creek valley, west of Picabo. To the east the northern limit runs across the Snake River Plain, and appears to be defined by increased effective precipitation and a general transformation to heavier soils, as evidenced by the increased presence of *Artemisia tripartita* and by the predominance of *Festuca idahoensis* as the understory dominant in the sagebrush-steppe, as opposed to the predominance of *Agropyron spicatum* and *Stipa comata* within its range. Superimposed on this precipitation and soil gradient are the recent lava flows between Carey and Laidlaw Park, which limit its distribution over large areas.

Western Limit: The western limit runs largely on a line between Picabo, Richfield and Shoshone, along lower Silver Creek and the Little Wood River. Most of the western edge of its distribution is defined by unsuitable soil types. To the west, no sandy loess occurs and the soils are high in clay. The result is the exclusive, but contiguous, distributions of two endemic milkvetch, *Astragalus atratus* var. *inseptus* and Picabo milkvetch. *Astragalus atratus* var. *inseptus* occurs on the clayey soils to the west and, with the exception of a small disjunct population of Picabo milkvetch (036), does not occur within the range of Picabo milkvetch (Smithman 1989; Prentice 1993). The disjunct population occurs on a small sandy deposit at the base of the Picabo Hills, approximately five miles east of the main portion of its range.

Extensive areas of sandy soil occur between Shoshone and Gooding, yet only two disjunct populations of Picabo milkvetch are known from this area. The distributional limit here, and to the south, may be defined by increased aridity, opposite of its northern limit.

Southern Limit: The southern limit runs northeast from Shoshone, across the Snake River Plain to just beyond the eastern edge of the Craters of the Moon Lava Field. Considerable suitable-appearing habitat occurs to the south, but only one disjunct population has been discovered southeast of Dietrich. As mentioned previously, increasing aridity to the south may limit its distribution here. Harrison (1981), Packard and Smithman (1984), and Popovich have searched extensively for Picabo milkvetch south of its known distribution.

Eastern Limit: The eastern limit of the main range of Picabo milkvetch is the Craters of the Moon Lava Field. Four disjunct occurrences, and a small area of suitable-appearing habitat, occur east of the lava field in the vicinity of Mule Butte. Beyond this small area, searches by Popovich (1994a) and, to a lesser extent, Moseley (1989) revealed little suitable habitat on the Big Desert, between the Craters of the Moon Lava Field and Big Southern Butte.

Extant occurrences: Appendix 3 contains the occurrence records for the 29 recently-visited occurrences of Picabo milkvetch (see below regarding 7 historical occurrences). These records contain detailed information on the location, population size, population area, habitat characteristics, occurrence documentation, and comments.

Extirpated occurrences: No populations of Picabo Milkvetch are known to be extirpated (but see discussion under historical occurrences, below).

Historical occurrences: Of the 36 documented occurrences in the CDC data base, 7 mentioned in Packard and Smithman's (1984) report have not been visited since 1984, and are considered historical. These include occurrences 005, 011, 012, 025, 027, 028, and 031 (Appendix 3). No population and habitat data has been documented for these sites and their current status is unknown. Occurrences 028 and 031 are among the most southern locations known.

Unverified/undocumented reports: A population of Picabo milkvetch in T55, R18E, S17, about four miles northeast of Shoshone, was reported by Packard and Smithman (1984) and is indicated on an undated, small-scale map on file at the Shoshone District BLM office. Authorship of the map is unknown. No other documentation exists for this population, and Moseley was unable to locate any Picabo milkvetch in section 17. A large depression surrounded by recent lava occurs in the center of the section, but no sandy soil was present. The only remotely potential habitat in this area is the small sand bars along the dry channel of the Big Wood River, a habitat that is probably too sandy to

be optimum habitat. We believe this is an erroneous occurrence and have not included it in the distribution of Picabo milkvetch (Appendices 2 and 3).

Synopsis of past and needed inventories: The distribution of Picabo milkvetch has been fairly well determined by numerous documented and undocumented surveys. Regarding documented surveys, Eidemiller found no milkvetch in her surveys of the foothills of the Pioneer, Smoky, and Soldier mountains (Eidemiller 1977a) and the Bennett Hills (Eidemiller 1977b). Rare plant inventories in the early 1980's contributed the most to our knowledge of the distribution of Picabo milkvetch west of the Craters of the Moon Lava Fields (Harrison 1981; Packard and Smithman 1984; Smithman 1984; Rosentreter 1986). Steve Popovich resumed surveying in the 1990's, with numerous searches in 1992, 1993, and 1994. A survey of sandy habitats east of the Craters of the Moon Lava Field by Popovich (1994a), resulted for the first time in the discovery of several populations on that side of the recent lavas. Popovich's 1994 survey, and field work by (Moseley 1989), confirmed that the appropriate sandy habitat required by Picabo milkvetch is common west of, but limited east of, these recent lava flows. Nearly all roads within the known range have been driven at least once in search of Picabo milkvetch. Many areas to the west, south, and southeast of the species known distribution have also been searched.

Little remains to be done as far as determining the general range of Picabo milkvetch. The boundaries of existing populations will certainly be expanded and new populations or subpopulations will probably be discovered within the range described above. New disjunct populations may be discovered as well. Searches of the kipukas in the lava between Laidlaw Park and Mule Butte should be conducted to determine if Picabo milkvetch occurs in these protected areas. Visits to historic sites should be made to determine their current status.

HABITAT

General habitat description: Picabo milkvetch occurs almost exclusively on the *Artemisia tridentata* var. *wyomingensis*/*Stipa comata* habitat type (Hironaka et al. 1983). This is an edaphically controlled habitat type restricted to sandy loam soils [in our case] or uniformly, highly calcareous silt loam soils. The sandy loam texture largely precludes *Agropyron spicatum* or *Stipa thurberiana* establishment. *Oryzopsis hymenoides* is a consistent member of this community (Hironaka et al. 1983). Topographically, Picabo milkvetch occupies sandy basins, bowls, and flats within rolling basalt. *Stipa comata* and *Oryzopsis hymenoides*, plants indicative of sandy soils in this area, are the best indicators of Picabo milkvetch habitat within the sagebrush/steppe. Dominant shrubs are *Artemisia tridentata* var. *wyomingensis*, *A. tridentata* subsp. *tridentata*, and *Chrysothamnus* spp. The precipitation zone is mostly 8-12", but it grades into the 12-16" zone along the northern edge of its distribution (Hironaka et al. 1983; Johnson 1991; Popovich 1994a). Slopes vary from nearly level to gently rolling. At these declivities, aspect is not a factor.

Elevations of known Picabo milkvetch populations range from 3700 feet for the disjunct populations between Shoshone and Gooding (028 and 034) to nearly 5200 feet at the northeastern limit of its distribution (007). The habitat at this latter site is considerably different than the others, being an *Artemisia tripartita*/*Agropyron spicatum* habitat (Hironaka et al. 1983) with an unusual assortment of associated species, such as *Carex filifolia*, *Koeleria cristata*, and *Phlox hoodii*. Sites 006, 008, and parts of 002 are ecotonal to *A. tripartita* habitat types, but are still dominated or co-dominated by *A. tridentata* and are not truly "clayey" sites.

As suggested earlier, the northern limit of its distribution is probably influenced by increased precipitation and increased clay content of the soils. *Artemisia tripartita* and *Festuca idahoensis* become more prominent beyond the northern limit of Picabo milkvetch. The southern limit appears to be influenced by hotter, more draughty conditions, and increased level of historic and current habitat alteration. Alteration consists of numerous, frequent wildfires, extensive invasion of weedy annuals, especially *Bromus tectorum*, high grazing levels, and mechanical seeding of *Agropyron cristatum*. At both the northern and southern limits, unobserved attributes, such as soil temperature and frost-free growing season, may also be affecting distribution.

Much of the species' occupied habitat, containing a majority of the known plants, burned in a series of large wildfires in 1992, which consumed 210,000 acres (Popovich 1994a). Picabo milkvetch plants appear to be more commonly found in open grassy areas that have been recently burned, than in closed-canopy sagebrush stands. Picabo milkvetch evolved with fire as part of the presettlement ecosystem processes and is able to survive burning. However, the fire regime has changed dramatically on the Snake River Plain within the last century, to the point where many native species are unable to withstand high fire frequencies and/or compete with cheatgrass (Pellant 1990; Whisenant 1990; Peters and Bunting 1994). Some sites containing very high densities (e.g., 017) are known to have burned at least 3-5 times since 1941. The impact of the this increased fire return interval on Picabo milkvetch is unknown. Fortunately, the sandy soils supporting Picabo milkvetch currently do not appear to be as prone to cheatgrass invasion as soils with siltier textures, although cheatgrass is very abundant in some localized sandy areas containing Picabo milkvetch.

Picabo milkvetch does not occupy unstable sands. *Psoralea lanceolata* characterizes these loose, shifting sands (Packard and Smithman 1984), and we never observed it to occur with Picabo milkvetch.

Geology and Soils: The bedrock underlying most of the range of Picabo milkvetch is basalt of the Snake River group, a common extrusive rock of Quaternary age that covers much of the eastern Snake River Plain. Undifferentiated Quaternary/Tertiary basalt underlies a few sites and Quaternary Wendell Grade basalt underlies the two occurrences (028 and 034) between Shoshone and Gooding (Rember and Bennett 1979a; 1979b; Worl et al. 1991). The few published K-Ar dates for the Snake River Group indicate that they were extruded less than 2 million years ago, although these dates have a large experimental error associated with them. Possibly a better indication is the normal polarity of exposed basalts, indicating that virtually all of them erupted in the last 750,000 years (Malde 1991). Following deposition of these basalts, sandy eolian loess was then deposited over much of the area, forming the parent material of the soils supporting Picabo milkvetch (Packard and Smithman 1984; Johnson 1991).

The western half of the range of Picabo milkvetch is interrupted by numerous Holocene lava flows of the Craters of the Moon Lava Field. Within the range of Picabo milkvetch, these flows range in age from approximately 12,000 years to less than 3600 years. Even younger flows occur to the north in Craters of the Moon National Monument. We speculate that these flows probably interrupted what was once a more continuous distribution of Picabo milkvetch. No loess was deposited on these recent flows and, therefore, they are unoccupied and unsuitable for Picabo milkvetch. The youngest flow within its range is the Minidoka Flow, which separates the eastern edge of the main distribution from the disjunct populations around Mule Butte. This flow, which separates these populations by nearly eight miles, is estimated to have been laid down 3590 ± 70 years ago (Kuntz et al. 1988).

The soil survey done for the Blaine County area (Johnson 1991), is the most recent soils mapping effort within the range of Picabo milkvetch. Although it does not cover the entire range, we will rely on this document to characterize the soils supporting the milkvetch. Three general groups of "well-drained soils on basalt plains" predominate:

1. McCary-Justesen-Rock Outcrop - These occur in the northeastern portion of its distribution in the Little Wood River valley. They are moderately deep and very deep soils that formed in loess over basalt residuum. These soils contain extensive and dense populations of Picabo milkvetch.
2. Deerhorn-Rehfield-Rock Outcrop - This general soil group occurs in the southern portion of Laidlaw Park and to the south of the recent flows and contains many, high density populations. They are moderately deep and very deep soils that formed in eolian material over basalt.
3. McCary-McBiggam-Bancroft - These soils occur in the Paddleford Flat, Little Park, and the northern portion of the Laidlaw Park kipukas. While these soils contain a few populations of Picabo milkvetch, they largely lie to the north of the core of its distribution. They are moderately deep and very deep soils that formed in loess over basalt residuum and in silty alluvium. Soil series in this group generally occur in higher precipitation zones than the previous two groups.

Refer to the soil survey (Johnson 1991) for a detailed description of each of the soil series.

Packard and Smithman (1984) noted that the species also occurs on clayey soils, as evidenced by the presence of *Viola beckwithii* and other clay-soil obligates. They didn't reference the location of this observation(s), but this situation has not been observed by other surveyors. In fact, Moseley observed that if there is enough clay in the soil to form polygonal cracks after drying, then Picabo milkvetch is always absent.

Associated species:

Shrubs - *Artemisia tridentata* var. *wyomingensis*, *A. tridentata* var. *tridentata*, *A. tripartita*, *Chrysothamnus nauseosus*, *C. viscidiflorus*, *Purshia tridentata*, *Tetradymia canescens*.

Graminoids - *Stipa comata*, *Oryzopsis hymenoides*, *Agropyron smithii/dasytachyum*(?), *A. spicatum*, *A. cristatum*, *Poa secunda*, *Bromus tectorum*, *Carex douglasii*, *C. filifolia*, *Koeleria cristata*.

Forbs - *Chorizanthe* sp., *Astragalus purshii*, *A. lentigenosus*, *Chaenactis douglasii*, *Sphaeralcea munroana*, *Gymnosteris nudicaulis*, *G. parvula*, *Cryptantha circumscissa*, *Alyssum desertorum*, *Eriogonum vimineum*, *Lomatium foeniculaceum*, *L. triternatum*, *Oenothera caespitosa*, *Lupinus* sp., *Phacelia heterophylla*, *Agoseris glauca*, *Phlox acaulis*, *P. longifolia*, *P. hoodii*, *Erigeron pumilus*, *Sisymbrium altissimum*, *Machaeranthera canescens*, *Leptodactylon pungens*, *Centaurea* sp., *Verbena* sp., *Penstemon acuminatum*, *Ranunculus glaberrimus*, *Opuntia polyacantha*, *Mimulus nanus*, *Collinsia parviflora*.

Other rare plant species: As previously mentioned, another rare milkvetch, *Astragalus atratus* var. *inseptus*, has a contiguous, but not sympatric, range with Picabo milkvetch (Smithman 1989; Prentice 1993). As discussed earlier, differences in soil texture appear to be the primary factor driving the distribution of these two species.

Gymnosteris parvula, a state monitor species (Idaho Native Plant Society 1994) was found to co-occur with Picabo milkvetch at one site near Mule Butte (033). The *Gymnosteris* occupied the more silty/clayey edge of the site while the greatest concentration of milkvetch occurred in the sandiest areas.

Moseley found *Mentzelia congesta*, a state Review species (Idaho Native Plant Society 1994), in unstable sands between Shoshone and Gooding, in the vicinity of Picabo milkvetch occurrences 028 and 034.

Gymnosteris nudicaulis occurs nearly throughout the range of Picabo milkvetch. This species was found to be more common than previously thought, and was recently dropped from the rare plant list for the state (Idaho Native Plant Society 1994). Although 1994 was a relatively dry spring, we found this annual to be widely distributed throughout the survey area.

POPULATION BIOLOGY

Phenology: Seed germination probably occurs early in the spring (or possibly in late fall). Flowering begins in mid-May most years. Fruit maturation proceeds through June, with most probably dehiscing sometime in July. There is wide variation in all these dates, possibly by as much as four weeks, depending on the temperature pattern during the spring.

Population size and condition: Appendix 3 contains the CDC data base records for the 36 known occurrences of Picabo milkvetch. These occurrence records contain information on location, survey dates, occurrence rank, population and habitat data, population size, area occupied, and various comment fields related to protection, management, and occurrence documentation. Below is a summary of selected occurrence record fields related to population size and condition. The Occurrence Rank is a relative ranking between A (highest) and D (lowest) based on population size, structure, and habitat quality. An Occurrence Rank of H refers to the seven historical collections that have not been revisited since they were discovered in 1984. Most of the population numbers and areas are estimates.

Occurrence Number	Occurrence Rank	Number Individuals	Area (acres)
001	C	100's	10+
002	B	1000's	500+
003	B	1000's	40+
004	C	100's	20
005	H	??	??
006	C	100's	5
007	B	1000's	10

Occurrence Number	Occurrence Rank	Number Individuals	Area (acres)
continued			
008	C	10	10?
009	A	10,000's	200+
010	A	10,000's	200+
011	H	??	??
012	H	??	??
013	A	10,000's	500+
014	A	1000	40
015	A	10,000's	500+
016	C	< 100	40
017	A	10,000's	hundreds
018	C	100	1
019	C	100's	40
020	C	100's	20?
021	D	100	3?
022	D	30	3?
023	A	1000's	40+
024	A	< 10,000	40
025	H	??	??
026	B	1000's	10
027	H	??	??
028	H	??	??
029	D	< 50	2
030	C	1000's	10
031	H	??	??
032	B	100	3
033	A	1100	11
034	D	200	100
035	D	200	100
036	C	100's	2

Steve Popovich, Shoshone BLM, and David Pyke, National Biological Service, began monitoring selected populations and habitats of Picabo milkvetch in 1993 and 1994, in response to wildfire and rehabilitation treatments, livestock grazing, and "range improvement plow and seed" treatment (Popovich 1994b). These monitoring sites are located at the following occurrences (see also Appendix 3 for site details):

1. Tikura plow and seed (occurrence 002)
2. Mile marker 186 on Hwy 26 (occurrence 017)
3. Pagari Bridge (occurrence 017)

4. Black Ridge Fire WSA site 2 (occurrence 014)
5. Black Ridge Fire WSA site 3 (occurrence 015)
6. Great Rift Fire 1 (occurrence 013)
7. Great Rift Fire 2 (occurrence 013)

Because the plots were only recently established, it is too early to discern population trends and all but tentative differences between treatments.

Reproductive Biology: Picabo milkvetch reproduces by seed. Nothing is known about seed dispersal, seed viability, or pollinators. Insects are assumed to be the pollination vector, therefore, it is possible that there is genetic interchange between all populations, even the disjunct ones. Genetic differentiation between populations has not been measured, however, to determine if this is the case. Popovich and Pyke have collected specimens of insects observed around plants at some monitoring sites, but they have not yet been identified.

Many Picabo milkvetch juveniles were observed in 1994. These plants probably germinated during 1993, which was an exceptionally wet spring and summer. We observed an enormous number of one-year old plants of many species of the sagebrush-steppe on the Snake River Plain.

The extent of the seed bank for this species is unknown, but it is widely known that many *Astragali* have seeds that remain viable for many years (Barneby 1964).

Biological Interactions: Harrison (1981) observed many plants to be chlorotic later in the growing season. He suspected that it was caused by insect predation, possibly a stem borer. He discounted other possibilities, such as mineral deficiency, because the symptoms would have been more widespread. Packard and Smithman (1984), however, suggested that drought is more likely the factor responsible. Popovich and Pyke observed that a small proportion of pods at their Tikura plow and seed study site had borers inside. One borer was captured, but they were unsuccessful in attempts to raise it to adult stage for identification.

Competition: It appears that Picabo milkvetch is not a good competitor. As stated previously, the sandy habitat occupied by Picabo milkvetch is often, but not always, very open, with considerable bare ground between plants. Newly open or disturbed habitat, such as roads, appears to be readily invaded by the milkvetch if the frequency of disturbance and competition is not too severe.

Herbivory: At most sites we did not observe significant herbivory of Picabo milkvetch plants in 1994. However, Popovich observed rabbit or small rodent herbivory to be 65 and 35 percent, respectively, for juvenile and adult plants at the Mile 186 monitoring site ($n = 20$ for each age class), and 29 and 11 percent, respectively, at the Tikura study site ($n = 35$ for each age class). When herbivory occurred, it was almost always severe, removing most or all above-ground phytomass. Sometimes some uneaten fruiting stalks remained nearby. Herbivory was not mentioned by any other botanists who conducted surveys for it prior to 1994.

Land ownership: Below is a table summarizing the ownership information of known Picabo milkvetch sites (see also Appendix 3). The BLM manages all or a portion of 32 occurrences (89%). The BLM shares management of these populations with the Idaho Department of Lands, and to a much lesser extent with the Idaho Department of Fish and Game and private owners. The Monument Resource Area, Shoshone District, has the greatest management responsibility with 26 occurrences.

The Bennett Hills Resource Area manages the two occurrences between Shoshone and Gooding, and the Idaho Falls District manages the four occurrences east of the Craters of the Moon Lava Field.

Occurrence Number	Ownership
001	Railroad and highway right-of way
002	BLM - Monument Resource Area
003	BLM - Monument Resource Area, State Lands
004	BLM - Monument Resource Area
005	BLM - Monument Resource Area
006	BLM - Monument Resource Area, Private
007	BLM - Monument Resource Area
008	BLM - Monument Resource Area
009	BLM - Monument Resource Area
010	BLM - Monument Resource Area, State Lands
011	State Lands
012	BLM - Monument Resource Area
013	BLM - Monument Resource Area, State Lands
014	BLM - Monument Resource Area
015	BLM - Monument Resource Area, State Lands
016	BLM - Monument Resource Area
017	BLM - Monument Resource Area, State Lands, Idaho Fish and Game, Private
018	State Lands
019	BLM - Monument Resource Area, State Lands
020	BLM - Monument Resource Area
021	BLM - Monument Resource Area
022	BLM - Monument Resource Area
023	BLM - Monument Resource Area, State Lands
024	BLM - Idaho Falls District
025	State Lands
026	BLM - Monument Resource Area, State Lands
027	BLM - Monument Resource Area
028	BLM - Bennett Hills Resource Area
029	BLM - Idaho Falls District
030	BLM - Monument Resource Area
031	BLM - Monument Resource Area
032	BLM - Idaho Falls District
033	BLM - Idaho Falls District
034	BLM - Bennett Hills Resource Area
035	BLM - Monument Resource Area
036	BLM - Monument Resource Area

Land use: The primary land use of virtually all habitat occupied by Picabo milkvetch is livestock grazing, both cattle and sheep. Portions of two occurrences do occur in areas that are ungrazed. These include a few plants in the fenced road right-of-way along lower Silver Creek (002) and a small portion of the population along the Little Wood River (017) that occurs in Bear Track Williams Wildlife Management Area, managed by the Idaho Department of Fish and Game.

ASSESSMENT AND MANAGEMENT RECOMMENDATIONS

Threats to currently known populations: The impacts to this plant from management activities are little understood. Observations have shown that populations may be present in areas receiving disturbance from light to moderate livestock grazing, invasion of annual exotics, increased fire frequency, and mechanical application of crested wheatgrass. However, we know nothing of the disturbance thresholds, long-term stability of the populations, and response to fire rehabilitation.

Given these uncertainties and the limited geographic range of Picabo milkvetch, a conservative approach to threat assessment and conservation planning is in order. Both short- and long-term threats to population and species viability of Picabo milkvetch exist. First a discussion of the short-term threats. There has already been considerable habitat lost to paved highways, gravel and dirt roads, powerline rights-of-way, canals, railroad rights-of-way, agricultural conversion, and livestock sacrifice areas (water developments, salting, bedding grounds, etc.). All of these could potentially be threats to populations or portions of populations in the future. These threats tend to be more localized in nature than the long-term ones, and more easily accounted for in land management planning. Also in need of consideration is the use of herbicides, especially in road rights-of-way, and noxious weed control programs.

Of greater concern is the 130-year decline of the sagebrush-steppe ecosystem on the Snake River Plain (Ertter and Moseley 1993; Noss et al. 1993) and its long-term effect on the persistence of rare plants, such as *Lepidium papilliferum* (Moseley 1994) and Picabo milkvetch. Although livestock grazing appears to have no immediate direct effect on population viability, in general past and possibly current grazing tends to cause detrimental changes in the composition, structure, and function in arid ecosystems (Tisdale et al. 1969; Yensen 1980; Tisdale and Hironaka 1981; Pellant 1990; Fleischer 1994). Picabo milkvetch habitat occupies areas conducive to habitat alteration and desirable for land treatment practices. It has had and is currently being converted to crested wheatgrass and other non-native perennials, invaded by noxious weeds, and invaded by exotic annuals that increase the fire return frequency to the point at which natives may be unable to persist. These problems are not as acute within the range of Picabo milkvetch as on the western Snake River Plain, but they are a cause for concern. The ecosystem- and population-level monitoring being conducted by the National Biological Service and Shoshone District BLM needs to be sensitive to habitat change due to the effects of management practices within the range of Picabo milkvetch.

Although Picabo milkvetch is sometimes found in crested wheatgrass seedings, the introduction of this perennial grass may be harmful in two ways: increased competition and the disturbance associated with the mechanical application of seed. As has been found with other rare plants in Idaho (Moseley 1994), plowing is probably more destructive to the habitat than broadcasting or drilling the seed. Packard and Smithman (1984) observed that the most continuous and highest density populations of Picabo milkvetch were indeed in crested wheatgrass seedings. Other surveyors, including both authors of this report, have observed the opposite. For example, near Brokie Lake (015) a sandy area

was partially plowed and there are noticeably fewer Picabo milkvetch plants on the disked area than in the adjacent undisturbed soil. Based on ocular estimates by Popovich in 1994, the one study site at Pagari that underwent a seeding contains less milkvetch than the three less-disturbed sites. Also, Popovich (unpublished data on file at the Shoshone District BLM office) found in 1994 that Picabo milkvetch was 6.5 times more abundant in an untreated sagebrush island as compared to an adjacent area that had been plowed and seeded in 1993. There was also a significant ($p < 0.05$) reduction in the proportion of seedlings and juveniles to adults in the treatment as compared to the untreated control plots. Unfortunately, pre-disturbance data is not available for any of these observations, limiting confidence in their interpretation.

Seeding exotic genotypes as part of a fire rehabilitation program should be critically evaluated. Following the Great Rift and Potter Butte fires in 1992, the natural sagebrush community in Laidlaw Park responded beautifully, with little mortality of forbs and grasses. The rehabilitation seeding that took place in this community was largely unnecessary to control erosion or prevent invasion of exotic annuals. If the pre-fire ecological status of the community is good, then little rehabilitation needs to be done (Peters and Bunting 1994). If, on the other hand, the ecological condition is poor and exotic annuals comprise a large part of the community, then seeding may be appropriate to prevent total conversion to annuals.

Diffuse knapweed (*Centaurea diffusa*) occurs in localized, but dense stands in disturbed areas around Picabo and Richfield. While not yet widespread in the area, it could pose a potential threat in the future if it increases in area and density.

Recommendations:

- o It is clear that the greatest threat to the viability of Picabo milkvetch is the continued decline of the native sagebrush-steppe on the Snake River Plain. The Shoshone District BLM should concentrate on this aspect of biodiversity conservation first and foremost. Picabo milkvetch is a narrowly-distributed endemic to this region, yet is locally abundant and not in such dire status that its immediate viability is in question. Any conservation strategy developed to assure the long-term persistence of Picabo milkvetch as a viable evolutionary unit must take into account the decline of the ecosystem.
- o In this ecosystem conservation context, we offer these general recommendations that should be considered when developing a conservation plan. While the largest populations, those ranked "A" and "B", are often thought to make the largest contribution to evolutionary viability, the importance of all extant populations to maintaining acceptable metapopulation structure and dynamics should not be discounted. A metapopulation is defined as a collection of "interdependent populations affected by recurrent extinctions and linked by recolonizations" (Murphy et al. 1990). In other words, while the loss of a few individuals may be deemed insignificant, it will generally be very important to maintain the overall geographic structure of the populations and their component subpopulations. The maintenance of the "distribution viability" of populations and subpopulations will serve as a good stand-in for maintaining less easily observed features that affect population and species viability, including genetic variation patterns, pollinator relationships, seed dispersal patterns, and gene flow within and among populations (Shelly 1994).

Maintaining metapopulation structure and dynamics becomes critical in the case of a narrow endemic like Picabo milkvetch, where many of the populations are isolated. Maintaining multiple populations over a wide range of geography and habitats will serve as a source for colonists as a hedge against a shifting mosaic of habitats and environmental stochasticity, and will buffer the effects of land management-induced alterations and future natural habitat changes (Shelly 1994).

- o Recommendation to the U.S. Fish and Wildlife Service: continue to maintain Picabo milkvetch as a category 3c candidate. If monitoring detects serious negative changes in ecosystem or population health, then it can be moved to the C1 category and possibly to listing, if appropriate.
- o Recommendation to the Idaho Native Plant Society: Continue to include Picabo milkvetch as a Monitor species, indicating that it is common within a limited range, but that it must continue to be monitored to detect possible upward or downward population trends.
- o Recommendation to the Idaho BLM: Continue to maintain Picabo milkvetch as a sensitive special status species.
- o Recommendations to the Shoshone and Idaho Falls Districts, BLM:
 - > Conduct a search of kipukas on the Minidoka Flow, between Mule Butte and Laidlaw Park, to determine if Picabo milkvetch occurs in these protected areas. Presence in the kipukas would be an important factor in developing conservation strategies.
 - > Relocate the seven historical sightings on the Shoshone District and assess their current status.
 - > Conduct intensive clearances of all proposed land development or land exchange projects that occur within the range outlined on the 1:100,000-scale maps. We know a considerable amount regarding the general distribution and habitat of Picabo milkvetch, but more intensive, local inventories will certainly extend the boundaries of known occurrences and possibly discover additional sites within the known range. There is also the possibility of new disjunct populations being discovered outside of this range.
 - > In order to maintain the "distribution viability" of Picabo milkvetch metapopulations, populations, and subpopulations, all habitat containing this narrow endemic should be given some form of special management consideration. A planning strategy that incorporates long-term species and habitat viability objectives in the BLM's broader land management planning process is highly recommended. This would include specific objectives whose achievement or failure would be measured by a monitoring program. A monitoring program would also ensure early detection of a downward trend. This would also include time-lines and clearly defined accountability. Above and beyond this, however, four areas deserve special recognition as ACEC/RNAs for their high-quality habitat and excellent populations of Picabo milkvetch:

Mule Butte - This site on the Idaho Falls District contains occurrence 024 (Appendix 2), a large dense population in high quality stands of the *Artemisia tridentata* var. *wyomingensis/Stipa comata* habitat type. The Mule Butte population is also important because it is the largest population on the eastern edge of its range, possibly having been isolated from the nearest populations for at least 3600 years by the Minidoka Flow. The area is relatively isolated and no permanent water sources have been developed nearby. The size of the special management area should encompass more than the known occupied area, perhaps a buffer of several miles on each side or incorporating the other sites nearby. An area of approximately 7500 acres will sufficiently encompass all known populations east of the lava flows and appropriate suitable habitat to maintain long-term metapopulation viability.

Bear Den Basin - Located in the eastern thumb of Laidlaw Park, in the basin between Bear Den Butte and Lava Butte, this site contains occurrence 009 (Appendix 2). Like the Mule Butte site, this isolated area has a large, high density population of Picabo milkvetch in a stand of *Artemisia tridentata* var. *wyomingensis/Stipa comata* in high ecological condition. The basin is traversed by a dirt road, but is isolated from most other developments. An area approximately 4500 acres in size will provide sufficient protection to this population and habitat.

Southeast of Pagari Bridge - The large sandy depression bisected by a two-track road in Section 6 (T4S R21E) of occurrence 017 (Appendix 2), about four miles southeast of Pagari Bridge, contains excellent habitat and high Picabo milkvetch densities in the northwestern portion of its range. There is also a permanent study plot there and several nearby along the road to Pagari Bridge. The entire area, from the bridge to Section 6, is being used to monitor long-term changes in plant populations and habitat attributes due to various land treatment practices.

Squaw Butte - The northern half of occurrence 015 (Appendix 2), from one mile south of Squaw Butte, along both sides of the road, several miles northwest toward The Blowout. This area is probably the best representation, in terms of habitat quality and plant numbers, of all known populations in the Wagon Butte to Black Ridge Crater area. This area encompasses the east-central portion of the species geographic range. Only the roadside has been surveyed; the full extent of the population is not known and may be considerably larger.

REFERENCES

- Barneby, R.C. 1957. *Pugillus Astragalorum* XX: Notes on *A. mulfordae* [sic] and some close relatives. Leaflets in Western Botany 8:120-125.
- Barneby, R.C. 1964. Atlas of North American *Astragalus*. Memoirs of the New York Botanical Garden Vol. 13. The New York Botanical Garden, Bronx, NY. 1188 p.
- Barneby, R.C. 1989. Fabales, Volume 3, Part B, Intermountain Flora. The New York Botanical Garden, Bronx, NY. 279 p.

- Conservation Data Center. 1994. Rare, threatened, and endangered plants and animals of Idaho. Third edition. Idaho Department of Fish and Game, Boise, ID. 39 p.
- Eidemiller, B.J. 1977a. Endangered and threatened plant inventory, Sun Valley ES area. Unpublished report prepared for the Shoshone District, BLM, Shoshone, ID. 33 p.
- Eidemiller, B.J. 1977b. Bennett Hills Environmental impact study area, threatened and endangered plant inventory report. Unpublished report prepared for the Shoshone District, BLM, Shoshone, ID. 56 p.
- Erter, B., and B. Moseley. 1992. Floristic regions of Idaho. *Journal of the Idaho Academy of Science* 28:57-70.
- Fleischner, T.L. 1994. Ecological costs of livestock grazing in western North America. *Conservation Biology* 8:629-644.
- Harrison, B.F. 1981. Inventory of threatened and endangered plants in a portion of the Shoshone District in Butte, Blaine, Lincoln, Minidoka, and Power counties, Idaho. Unpublished report on file at the Shoshone District, BLM, Shoshone, ID. 27 p. plus appendix.
- Hironaka, M., M.A. Fosberg, and A.H. Winward. 1983. Sagebrush-grass habitat types of southern Idaho. Bulletin Number 35. Forest, Wildlife and Range Experiment Station, University of Idaho, Moscow, ID. 44 p.
- Hitchcock, C.L. 1961. *Astragalus*. Pages 197-273 in: *Vascular plants of the Pacific Northwest, Part 3*, by C.L. Hitchcock, A. Cronquist, M. Ownbey, and J.W. Thompson. University of Washington Press, Seattle.
- Hitchcock, C.L., and A. Cronquist. 1973. *Flora of the Pacific Northwest*. University of Washington Press, Seattle, WA. 730 p.
- Idaho Native Plant Society. 1994. Results of the tenth annual Idaho Rare Plant Conference. Unpublished document on file at the Conservation Data Center, Idaho Department of Fish and Game, Boise, ID.
- Isely, D. 1986. Leguminosae of the United States. *Astragalus* L.:IV. Species summary N-Z. *Iowa State Journal of Research* 62:157-289.
- Johnson, M.E. 1991. Soil survey of Blaine County area, Idaho. USDA, Soil Conservation Service, Boise, ID. 388 p.
- Johnson, F.D., and R.W. Steele. 1974. A tentative list of uncommon plants of Idaho. Pages 105-123 in: *Research Natural Needs in Idaho*, C.A. Wellner and F.D. Johnson, eds., College of Forestry, Wildlife and Ranges Sciences, University of Idaho, Moscow, ID.

- Kuntz, M. D.E. Champion, R.H. Lefebvre, and H.R. Covington. Geologic map of the Craters of the Moon, Kings Bowl, and Wapi lava fields, and the Great Rift Volcanic Rift Zone, south-central Idaho. Miscellaneous Investigation Series Map I-1632. U.S.D.I., Geological Survey, Denver, CO.
- Malde, H.E. 1991. Quaternary geology and structural history of the Snake River Plain, Idaho and Oregon. Pages 251-281 in *Quaternary Nonglacial Geology: Conterminous U.S.*, The Geology of North America, Vol. K-2, R.B. Morrison, ed., Geological Society of America, Boulder, Co.
- Moseley, R.K. 1989. Report of the conservation status of *Phacelia inconspicua* in Idaho. Unpublished report on file at the Conservation Data Center, Idaho Department of Fish and Game, Boise, ID. 18 p. plus appendices.
- Moseley, R.K. 1994. Report on the conservation status of *Lepidium papilliferum*. Unpublished report on file at the Conservation Data Center, Idaho Department of Fish and Game, Boise, ID. 35 p. plus appendices.
- Murphy, D.D., K.E. Freas, and S.B. Weiss. 1990. An environment-metapopulation approach to population viability analysis for a threatened invertebrate. *Conservation Biology* 4:41-51.
- Noss, R.F., E.T. LaRoe, and J.M. Scott. 1993. Endangered ecosystems of the United States. A preliminary assessment of loss and degradation. Draft manuscript on file at the Idaho Department of Fish and Game, Conservation Data Center, Boise, ID. 71 p.
- Packard, P.L. 1981. *Astragalus oniciformis*. Page 8 in: *Vascular plant species of concern in Idaho, by the Rare and Endangered Plants Technical Committee of the Idaho Natural Areas Council*. Bulletin Number 34, Forest, Wildlife and Range Experiment Station, University of Idaho, Moscow, ID.
- Packard, P.L., and L.C. Smithman. 1984. Threatened and endangered species training and inventory project, *Astragalus oniciformis* Barneby. Unpublished report on file at the Shoshone District, BLM, Shoshone, ID. 17 p. plus appendices.
- Pellant, M. 1990. The cheatgrass-wildfire cycle - Are there any solutions? Pages 11-18 in: *Proceedings-Symposium on Cheatgrass Invasion, Shrub Die-off, and Other Aspects of Shrub Biology and Management*, E.D. McArthur, E.M. Romney, S.D. Smith, and P.T. Tueller, compilers, General Technical Report INT-276, USDA Forest Service, Intermountain Research Station, Ogden, UT.
- Peters, E.F., and S.C. Bunting. 1994. Fire conditions pre- and postoccurrence of annual grasses on the Snake River Plain. Pages 31-36 in: *Proceedings-Ecology and Management of Annual Rangelands*, S.B. Monsen and S.G. Kitchen, eds., General Technical Report INT-GTR-13, USDA Forest Service, Intermountain Research Station, Ogden, UT.
- Popovich, S.J. 1994a. Field survey for *Astragalus oniciformis* along the eastern edge of the Great Rift lava fields. Unpublished report on file at the Shoshone District, BLM, Shoshone, ID. 5 p. plus maps.


- Popovich, S.J. 1994b. *Astragalus oniciformis* study site and methodology information. Unpublished report on file at the Shoshone District, BLM, Shoshone, ID.
- Prentice, C.A. 1993. Camas milkvetch inventory, *Astragalus atratus* S.Wats. var. *inseptus* Barneby. Unpublished report on file at the Shoshone District BLM, Shoshone, ID. 35 p.
- Rember, W.C., and E.H. Bennett. 1979a. Geologic map of the Idaho Falls Quadrangle, Idaho. Geologic Map Series (2nd Quadrangle). Idaho Bureau of Mines and Geology, Moscow, ID
- Rember, W.C., and E.H. Bennett. 1979b. Geologic map of the Twin Falls Quadrangle, Idaho. Geologic Map Series (2nd Quadrangle). Idaho Bureau of Mines and Geology, Moscow, ID
- Rosentreter, R. 1986. Sensitive and uncommon plants in the Shoshone District, Bureau of Land Management, 1985. Unpublished report on file at Idaho State Office, BLM, Boise, ID. 23 p.
- Shelly, S. 1994. Population viability assessments for TES plants. Draft. Unpublished report prepared for the U.S. Forest Service, on file at the Idaho Department of Fish and Game, Conservation Data Center, Boise, ID.
- Smithman, L.C. 1984. Threatened and endangered plant survey, proposed Silver-Midpoint Transmission Line. Unpublished report prepared for Idaho Power Company, Boise, ID. 15 p.
- Smithman, L.C. 1989. Threatened and endangered species training and inventory project: *Astragalus atratus* Wats. variety *inseptus* Barneby. Technical Bulletin 89-1. Idaho State Office, Bureau of Land Management, Boise, ID 22 p.
- Steele, R.W. 1975. A directory of disjunct and endemic plants of central and southern Idaho. Information Series Number 9. Forest, Wildlife and Range Experiment Station, University of Idaho, Moscow, ID. 26 p.
- Tisdale, E.W., M. Hironaka, and M.A. Fosberg. 1969. The sagebrush region of Idaho. A problem in range resource management. Bulletin 512. Agricultural Experiment Station, University of Idaho. 15 p.
- Tisdale, E.W., and M. Hironaka. 1981. The sagebrush-grass region: A review of the ecological literature. Bulletin Number 33. Forest, Wildlife and Range Experiment Station, University of Idaho. 31 p.
- U.S. Fish and Wildlife Service. 1980. Endangered and threatened wildlife and plants; review of plant taxa for listing as endangered or threatened species. Federal Register 50 CFR Part 17 45(242):82480-82569 (December 15, 1980).
- U.S. Fish and Wildlife Service. 1983. Endangered and threatened wildlife and plants; supplement to review of plant taxa for listing; proposed rule. Federal Register 50 CFR Part 17 48(229):53640-53670 (November 28, 1983).

- U.S. Fish and Wildlife Service. 1985. Endangered and threatened wildlife and plants; review of plant taxa for listing as endangered or threatened species; notice of review. Federal Register 50 CFR Part 17 50(188):39526-39527 (September 27, 1985).
- U.S. Fish and Wildlife Service. 1990. Endangered and threatened wildlife and plants; review of plant taxa for listing as endangered or threatened species; notice of review. Federal Register 50 CFR Part 17 55(35):6184-6229 (February 21, 1990).
- U.S. Fish and Wildlife Service. 1993. Plant taxa for listing as endangered or threatened species; notice of review. Federal Register 50 CFR Part 17 58(188):51144-51190 (September 30, 1993).
- Wiseman, S.G. 1990. Changing fire frequencies on Idaho's Snake River Plains: Ecological and management implications. Pages 4-10 in: Proceedings-Symposium on Cheatgrass Invasion, Shrub Die-off, and Other Aspects of Shrub Biology and Management, E.D. McArthur, E.M. Romney, S.D. Smith, and P.T. Tueller, compilers. General Technical Report INT-276, USDA Forest Service, Intermountain Research Station, Ogden, UT.
- Worl, R.G., T.H. Kiilsgaard, E.H. Bennett, P.K. Link, R.S. Lewis, V.E. Mitchell, K.M. Johnson, and L.D. Snyder. 1991. Geologic map of the Hailey 1°x2° Quadrangle, Idaho. Open-file Report 91-340. USDI Geological Survey and Idaho Geological Survey, Moscow, ID.
- Yensen, D. 1980. A grazing history of southwestern Idaho with emphasis on the Birds of Prey Study Area. USDI-BLM Snake River Birds of Prey Research Project, Boise, ID. 82 p.
- Young, J.A. 1994. History and use of semiarid plant communities - changes in vegetation. Pages 5-8 in: Proceedings-Ecology and Management of Annual Rangelands, S.B. Monsen and S.G. Kitchen, eds., General Technical Report INT-GTR-13, USDA Forest Service, Intermountain Research Station, Ogden, UT.

Appendix 1

Line drawings of *Astragalus oniciformis*
(from Barneby 1989).

plant generally prostrate,
2-6 inches tall, up to 2
feet in diameter




Astragalus oniciformis
Picabo milkvetch



Appendix 2

Maps of the distribution of *Astragalus oniciformis*.



Distribution of *Astragalus oniciformis*
(note four outlying populations)




Distribution of *Astragalus oniciformis*
January, 1995

The three-digit code associated with the circled area occupied by each population or groups of populations is the occurrence number from the Idaho Conservation Data Center. Heavy dashed line denotes the approximate limit of suitable habitat. Photocopied and reduced to 60% of actual size from the 1990 Twin Falls 1:100,000-scale Idaho Transportation Department topographic map. Original map on file at the BLM-Shoshone District Office and the ICDC Boise Office.





Distribution of *Astragalus oniciformis*
January, 1995

The three-digit code associated with the circled area occupied by each population or groups of populations is the occurrence number from the Idaho Conservation Data Center. Heavy dashed line denotes the approximate limit of suitable habitat. Photocopied and reduced to 60% of actual size from the 1992 Fairfield 1:100,000-scale Idaho Transportation Department topographic map. Original map on file at the BLM-Shoshone District Office and the ICDC Boise Office.



Distribution of *Astragalus oniciformis*
January, 1995

The three-digit code associated with the circled area occupied by each population or groups of populations is the occurrence number from the Idaho Conservation Data Center. Heavy dashed line denotes the approximate limit of suitable habitat. Photocopied and reduced to 60% of actual size from the 1992 Fairfield 1:100,000-scale Idaho Transportation Department topographic map. Original map on file at the BLM-Shoshone District Office and the ICDC Boise Office.




Topographic Map Symbols

- Primary and Secondary highways
- Light duty road or road under const. or not open for traffic
- Other road or street type
- International border
- Service boundary of another country
- Service boundary of another state
- Service boundary of another territory
- Service boundary of another district
- Service boundary of another county
- Service boundary of another township
- Service boundary of another section
- Service boundary of another quarter section
- Service boundary of another acre
- Service boundary of another foot
- Service boundary of another inch
- Service boundary of another cent
- Service boundary of another mill
- Service boundary of another micron
- Service boundary of another nanometer
- Service boundary of another micrometer
- Service boundary of another millimeter
- Service boundary of another centimeter
- Service boundary of another decimeter
- Service boundary of another meter
- Service boundary of another kilometer
- Service boundary of another megameter
- Service boundary of another gigameter
- Service boundary of another terameter
- Service boundary of another petameter
- Service boundary of another exameter
- Service boundary of another zettameter
- Service boundary of another yottameter
- Service boundary of another ronnometer
- Service boundary of another quectometer
- Service boundary of another sexagesimal minute
- Service boundary of another sexagesimal second
- Service boundary of another sexagesimal third
- Service boundary of another sexagesimal fourth
- Service boundary of another sexagesimal fifth
- Service boundary of another sexagesimal sixth
- Service boundary of another sexagesimal seventh
- Service boundary of another sexagesimal eighth
- Service boundary of another sexagesimal ninth
- Service boundary of another sexagesimal tenth
- Service boundary of another sexagesimal eleventh
- Service boundary of another sexagesimal twelfth
- Service boundary of another sexagesimal thirteenth
- Service boundary of another sexagesimal fourteenth
- Service boundary of another sexagesimal fifteenth
- Service boundary of another sexagesimal sixteenth
- Service boundary of another sexagesimal seventeenth
- Service boundary of another sexagesimal eighteenth
- Service boundary of another sexagesimal nineteenth
- Service boundary of another sexagesimal twentieth
- Service boundary of another sexagesimal twenty-first
- Service boundary of another sexagesimal twenty-second
- Service boundary of another sexagesimal twenty-third
- Service boundary of another sexagesimal twenty-fourth
- Service boundary of another sexagesimal twenty-fifth
- Service boundary of another sexagesimal twenty-sixth
- Service boundary of another sexagesimal twenty-seventh
- Service boundary of another sexagesimal twenty-eighth
- Service boundary of another sexagesimal twenty-ninth
- Service boundary of another sexagesimal thirtieth
- Service boundary of another sexagesimal thirty-first
- Service boundary of another sexagesimal thirty-second
- Service boundary of another sexagesimal thirty-third
- Service boundary of another sexagesimal thirty-fourth
- Service boundary of another sexagesimal thirty-fifth
- Service boundary of another sexagesimal thirty-sixth
- Service boundary of another sexagesimal thirty-seventh
- Service boundary of another sexagesimal thirty-eighth
- Service boundary of another sexagesimal thirty-ninth
- Service boundary of another sexagesimal fortieth
- Service boundary of another sexagesimal forty-first
- Service boundary of another sexagesimal forty-second
- Service boundary of another sexagesimal forty-third
- Service boundary of another sexagesimal forty-fourth
- Service boundary of another sexagesimal forty-fifth
- Service boundary of another sexagesimal forty-sixth
- Service boundary of another sexagesimal forty-seventh
- Service boundary of another sexagesimal forty-eighth
- Service boundary of another sexagesimal forty-ninth
- Service boundary of another sexagesimal fiftieth
- Service boundary of another sexagesimal fifty-first
- Service boundary of another sexagesimal fifty-second
- Service boundary of another sexagesimal fifty-third
- Service boundary of another sexagesimal fifty-fourth
- Service boundary of another sexagesimal fifty-fifth
- Service boundary of another sexagesimal fifty-sixth
- Service boundary of another sexagesimal fifty-seventh
- Service boundary of another sexagesimal fifty-eighth
- Service boundary of another sexagesimal fifty-ninth
- Service boundary of another sexagesimal sixtieth
- Service boundary of another sexagesimal sixty-first
- Service boundary of another sexagesimal sixty-second
- Service boundary of another sexagesimal sixty-third
- Service boundary of another sexagesimal sixty-fourth
- Service boundary of another sexagesimal sixty-fifth
- Service boundary of another sexagesimal sixty-sixth
- Service boundary of another sexagesimal sixty-seventh
- Service boundary of another sexagesimal sixty-eighth
- Service boundary of another sexagesimal sixty-ninth
- Service boundary of another sexagesimal seventieth
- Service boundary of another sexagesimal seventy-first
- Service boundary of another sexagesimal seventy-second
- Service boundary of another sexagesimal seventy-third
- Service boundary of another sexagesimal seventy-fourth
- Service boundary of another sexagesimal seventy-fifth
- Service boundary of another sexagesimal seventy-sixth
- Service boundary of another sexagesimal seventy-seventh
- Service boundary of another sexagesimal seventy-eighth
- Service boundary of another sexagesimal seventy-ninth
- Service boundary of another sexagesimal eightieth
- Service boundary of another sexagesimal eighty-first
- Service boundary of another sexagesimal eighty-second
- Service boundary of another sexagesimal eighty-third
- Service boundary of another sexagesimal eighty-fourth
- Service boundary of another sexagesimal eighty-fifth
- Service boundary of another sexagesimal eighty-sixth
- Service boundary of another sexagesimal eighty-seventh
- Service boundary of another sexagesimal eighty-eighth
- Service boundary of another sexagesimal eighty-ninth
- Service boundary of another sexagesimal ninetieth
- Service boundary of another sexagesimal ninety-first
- Service boundary of another sexagesimal ninety-second
- Service boundary of another sexagesimal ninety-third
- Service boundary of another sexagesimal ninety-fourth
- Service boundary of another sexagesimal ninety-fifth
- Service boundary of another sexagesimal ninety-sixth
- Service boundary of another sexagesimal ninety-seventh
- Service boundary of another sexagesimal ninety-eighth
- Service boundary of another sexagesimal ninety-ninth
- Service boundary of another sexagesimal hundredth


Distribution of *Astragalus oniciformis*
January, 1995

The three-digit code associated with the circled area occupied by each population or groups of populations is the occurrence number from the Idaho Conservation Data Center. Heavy dashed line denotes the approximate limit of suitable habitat. Photocopied and reduced to 60% of actual size from the 1990 Twin Falls 1:100,000-scale Idaho Transportation Department topographic map. Original map on file at the BLM-Shoshone District Office and the ICDC Boise Office.




Distribution of *Astragalus oniciformis*
January, 1995

The three-digit code associated with the circled area occupied by each population or groups of populations is the occurrence number from the Idaho Conservation Data Center. Heavy dashed line denotes the approximate limit of suitable habitat. Photocopied and reduced to 60% of actual size from the 1991 Craters of the Moon 1:100,000-scale Idaho Transportation Department topographic map. Original map on file at the BLM-Shoshone District Office and the ICDC Boise Office.



Distribution of *Astragalus onictiformis*
January, 1995

The three-digit code associated with the circled area occupied by each population or groups of populations is the occurrence number from the Idaho Conservation Data Center. Heavy dashed line denotes the approximate limit of suitable habitat. Photocopied and reduced to 60% of actual size from the 1991 Craters of the Moon 1:100,000-scale Idaho Transportation Department topographic map. Original map on file at the BLM-Shoshone District Office and the ICDC Boise Office.



Distribution of *Astragalus oniciformis*
January, 1995

The three-digit code associated with the circled area occupied by each population or groups of populations is the occurrence number from the Idaho Conservation Data Center. Heavy dashed line denotes the approximate limit of suitable habitat. Photocopied and reduced to 60% of actual size from the 1991 Craters of the Moon 1:100,000-scale Idaho Transportation Department topographic map. Original map on file at the BLM-Shoshone District Office and the ICDC Boise Office.



Appendix 3

Occurrence records from the Conservation
Data Center for *Astragalus oniciformis*.

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 001

Survey Site Name: PICABO

County: Blaine

USGS quadrangle: PICABO

Latitude: 43 18 00 N Longitude: 114 02 20 W

TOWNRANGE:	SECTION:	TRSNOTE:
001S020E	26	SE4SE4SW4, SW4SW4SE4
001S020E	35	NE4, NE4NE4NW4
001S020E	36	SW4SW4NW4

Location:

Along old railroad right-of-way and US Route 20 right-of-way (south side), beginning about 1 mile east of Picabo store and continuing SE for ca 0.8 mile.

Survey Date: 1994-05-12

Last Observed: 1994-05-12

First Observed: 1941

EORANK: C

EORANK Comments: Fragmented and disturbed habitat.

Population Data:

1963: Corolla nearly white, the banner purple-lined, pods pendulous, triquetrous, 2-celled, plants prostrate. 1994: Several hundred plants in full flower; good age class distribution. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Narrow strip of habitat south of railroad right-of-way and between railroad and US Route 20. Sandy loam soil over basalt. Degraded *Artemisia tridentata wyomingensis*/*Stipa comata* habitat type with *Tetradymia canescens*, *Gymnosteris nudicaulis*, *Chrysothamnus nauseosus*, and several weeds including *Bromus tectorum*, *Centaurea* sp., and *Sisymbrium altissimum*.

Minimum Elevation: 4820 feet

Size: 10+ AC (LINEAR)

Ownership Comments:

Private land and/or highway right-of-way.

Comments:

Occurrence was compiled from the following sources: Packard and Smithman 1984 (voucher specimen indicated); small scale map (1981-1984?) in Shoshone District BLM Office; and Moseley 1994 survey.

Specimens:

R. Rosentreter 3641 (ID, MONTU, CIC).

Lynda C. Smithman LS-1121 (CIC).

Wood s.n. (IDS).

Ripley and Barneby 8795 (WTU, isotope; IDS, isotope; ID; IDF).

C. L. Hitchcock 22704 (ID).

C. V. Muhlick s.n. (ID).

R.J. Davis 3088 (IDS)

Nelson and Macbride (2987 (CAS, GH, MO, NY)

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 002

Survey Site Name: SILVER CREEK

County: Blaine

USGS quadrangle: PICABO
CAREY
PAGARI WELL

Latitude: 43 15 37 N Longitude: 114 00 47 W

TOWNRANGE:	SECTION:	TRSNOTE:
002S020E	01	
002S020E	12	E2, E2NW4
002S020E	13	NE4NE4
002S021E	07	W2, W2SE4
002S021E	08	
002S021E	17	N2NE4, NE4NE4NW4
002S021E	18	

Location:

Along Silver Creek upstream of US Route 20 at the BLM Silver Creek Access site and extending N in the Picabo Cutoff Road right-of-way. A nearby population is located E of Silver Creek, SE of Picabo.

Survey Date: 1994-05-12
Last Observed: 1994-05-12
First Observed: 1983-06-28

EORANK: B

EORANK Comments:

Population Data:

1994: Many thousands of individuals. Plants are widely scattered in the road right-of-way, but probably extend onto adjacent land in some cases. The populations on BLM land are moderately dense, although density decreases in crested wheatgrass seedings. Plants in full flower. Area surveyed by Bob Moseley, Idaho CDC. 1994 (Fall): One subpopulation thoroughly surveyed by S. Popovich, Shoshone District BLM, and David Pyke, NBS. 300 genets estimated, population vigor assessed as good.

Habitat Description:

Most of occurrence is in *Artemisia tridentata wyomingensis/Stipa comata* habitat type, much of it in degraded or altered condition. Many weeds in road right-of-way. Much of the area has been seeded to *Agropyron cristatum*. Some of the area was plowed prior to seeding. The remaining sagebrush stands are heavily grazed. Deep sandy loam over basalt with considerable bare ground.

Minimum Elevation: 4700 feet

Maximum Elevation: 4810 feet

Size: 500+ AC

Ownership Comments:

Shoshone District BLM, Monument RA; private land; and county road right-of-way.

Comments:

The occurrence was probably once continuous with *A. oniciformis* .001 to the north, but the intervening area is now irrigated cropland. The exact boundaries of the occurrence are unknown; it probably extends onto private land outside of the right-of-way. The occurrence was compiled from the following sources: Packard and Smithman 1984 (voucher specimen indicated), small-scale (1981-1984?) map at the Shoshone District BLM Office, Steve Popovich (Shoshone District BLM) 1993-1994 surveys, and Moseley 1994 survey.

Protection Comments:

Currently there is cattle grazing; possible future threats may be plow and seeding.

Management Comments:

Monitoring plots established in T2SR20E Section 1 (NW4SW4).

Specimens:

Lynda C. Smithman LS-116 (CIC).

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 003

Survey Site Name: PADDELFORD FLAT

County: Blaine

USGS quadrangle: PADDELFORD FLAT

Latitude: 43 17 25 N Longitude: 113 47 31 W

TOWNRANGE:	SECTION:	TRSNOTE:
001S022E	35	SE4SE4
001S022E	36	
002S022E	01	N2NW4
002S022E	02	NE4NE4

Location:

1.6 miles S of Paddleford Flat Lake; about 5 miles NE of Carey, turn E off Hwy 20/26/93 and proceed about 5 miles. Then turn S about proceed for about 2 miles.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1981-06-12

EORANK:

EORANK Comments:

Population Data: B

1994: Moderately dense population with good age class distribution; thousands of plants in full flower. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Unburned *Artemisia tridentata wyomingensis*/*Stipa comata* community on deep sandy loam soil (loess) over basalt. Level to gently sloping. Associated with *Bromus tectorum*, *Carex douglasii*, *Poa secunda*, and *Phacelia heterophylla*. Area is not heavily grazed - plant community is in good condition.

Minimum Elevation: 4800 feet

Maximum Elevation: 4820 feet

Size: 40 + AC

Ownership Comments:

State land and Shoshone District BLM, Monument RA.

Comments:

The boundaries of the population are not precisely known. This occurrence was compiled from the following sources: Harrison 1981, Packard and Smithman 1984 (voucher specimen indicated), small-scale (1981-1984?) map in the Shoshone District BLM Office, and Moseley 1994 survey.

Specimens:

B. F. and L. C. Harrison 13,447; 13,476 (Shoshone BLM).

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 004

Survey Site Name: WAGON BUTTE

County: Blaine

USGS quadrangle: WAGON BUTTE

Latitude: 43 13 18 N Longitude: 113 47 24 W

TOWNRANGE: SECTION: TRSNOTE:
002S022E 25 NE4SW4, SE4NW4

Location:

About 20 mi SE of Carey, E side of Wagon Butte, S of Paddleford Flat.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1984

EORANK: C

EORANK Comments:

Population Data:

1994: Hundreds of plants. Population is of moderate density with good age class distribution. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Unburned *Artemisia tridentata wyomingensis*/*Stipa comata* community in fair condition. High cover of *Bromus tectorum* and *Chrysothamnus viscidiflorus*. Deep sandy soil.

Minimum Elevation: 4720 feet

Maximum Elevation: 4740 feet

Size: 20

Ownership Comments: Shoshone District BLM, Monument RA.

Comments: The boundaries of the population are not precisely known. This occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale (1981-1984?) map in the Shoshone District BLM Office, 7.5' quad map (1981-1984?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Protection Comments: Water developments nearby have resulted in heavy cattle use.

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 005

Survey Site Name: INDIAN WELL

County: Blaine

USGS quadrangle: LITTLE PARK

Latitude: 43 15 48 N Longitude: 113 39 16 W

TOWNRANGE:	SECTION:	TRSNOTE:
002S023E	12	E2SE4
002S024E	07	W2SW4

Location:

About 19 mi E of Carey. Eastern edge of Laidlaw Park, S of North Entrance Indian Well along the road from Little Park.

Survey Date: 1994-05-11

Last Observed: 1984

First Observed: 1984

EORANK: H

EORANK Comments:

Population Data:

1994: Not relocated by Bob Moseley, Idaho CDC, during his 1994 surveys.

Habitat Description:

Unburned *Artemisia tridentata wyomingensis* and *Artemisia tripartita* communities.

Minimum Elevation: 4980 feet

Maximum Elevation: 5020 feet

Size:

Ownership Comments:

Shoshone District BLM, Monument RA.

Comments:

This occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 006

Survey Site Name: TWIN LAKES

County: Blaine

USGS quadrangle: BEAR DEN BUTTE
LAIDLAW BUTTE
NORTH LAIDLAW BUTTE

Latitude: 43 14 52 N Longitude: 113 37 37 W

TOWNRANGE:	SECTION:	TRSNOTE:
002S024E	08	NE4
002S024E	17	SW4
002S024E	20	NE4NE4NW4

Location:

About 21.5 mi SE of Carey. NE of Laidlaw Butte, W of Turnbull Butte, and N of Corral Butte.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1984-06-13

EORANK: C

EORANK Comments:

Population Data:

1993: Three mature flowering plants in one of the northern subpopulations was reported by Steve Popovich, Shoshone District BLM. Vigor assessed as good on this cursory visit. 1994: The northern portion of the occurrence (in section 8) was not relocated. East of Twin Lakes (section 17): moderately dense population with good age class distribution. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

The north end of the population burned in the 1980's (?). Unburned *Artemisia tridentata tridentata*/*Stipa comata* and *Artemisia tridentata wyomingensis*/*Stipa comata* communities near Twin Lakes. Sandy loam over basalt. Gentle slopes with considerable bare ground. Associated with *Chrysothamnus nauseosus*, *Agropyron smithii*, *A. spicatum*, *Poa secunda*, *P. sandbergii*, *Ranunculus glaberrimus*, and some *Artemisia tripartita*.

Minimum Elevation: 4970 feet
Maximum Elevation: 5100 feet
Size: 5 AC

Ownership Comments:
Shoshone District BLM, Monument RA, and private land.

Comments:
The precise boundaries of the population are unknown. Packard and Smithman (1984) indicates that the population occurs in sections 8 and 17. The occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale map (1981-1984?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Protection Comments:
Sheep have been grazing in the nearby area.

Specimens:
Lynda C. Smithman LS-1416 (ID, CIC).

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 007

Survey Site Name: BULLSHOT RESERVOIR

County: Blaine

USGS quadrangle: NORTH LAIDLAW BUTTE

Latitude: 43 16 13 N Longitude: 113 33 12 W

TOWNRANGE:	SECTION:	TRSNOTE:
002S024E	11	NE4NE4
002S024E	12	NW4NW4

Location:

1.4 Air miles NE of Turnbull Butte and ca 1.6 road miles SW of Bullshot Reservoir.

Survey Date: 1994-05-11
Last Observed: 1994-05-11
First Observed: 1984

EORANK: B
EORANK Comments:

Population Data:

1994: Thousands of individuals. Population has good age class distribution. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Unburned *Artemisia tripartita*/*Agropyron spicatum* community on level sandy soil. Considerable bare ground. Associated species include *Phlox hoodii*, *Gymnosteris nudicaulis*, *Carex douglasii*, *Stipa comata*, *Poa secunda*, *Carex filifolia*, *Opuntia polyacantha*, and *Koeleria cristata*.

Minimum Elevation: 5160 feet

Size: 10

Ownership Comments:

Shoshone District BLM, Monument RA: population might extend on to private land.

Comments:

The precise boundaries of the population are unknown. The occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale map (1981-1984?) in the Shoshone District BLM Office, 7.5' quad in the Shoshone District BLM Office, and Moseley 1994 survey.

Specimens:

· *ASTRAGALUS ONICIFORMIS*

PICABO MILKVETCH

Occurrence Number: 008

Survey Site Name: CREAM CAN LAKE

County: Blaine
Minidoka

USGS quadrangle: BEAR DEN BUTTE

Latitude: 43 12 07 N Longitude: 113 36 22 W

TOWNRANGE:	SECTION:	TRSNOTE:
002S024E	28	SW4SW4NE4, NW4NW4SE4
002S024E	33	SW4, W2W2SE4
003S024E	04	N2N2NW4

Location:
Ca 24 and 25 miles SE of Carey along and north of the Blaine
County/Minidoka County line. About 1 to 1.75 miles north of Cream
Can Lake.

Survey Date: 1994-05-11
Last Observed: 1994-05-11
First Observed: 1981-06-11

EORANK: C
EORANK Comments:

Population Data:
1993-1994: The northern population contains about 10 scattered
plants. Observation by Steve Popovich, Shoshone District BLM. The
area was not visited by Bob Moseley, Idaho CDC, in 1994. 1994: 10
genets observed on this cursory visit by S. Popovich. Population
assessed as fair, and no change since last visit.

Habitat Description:
Sandy clay loams over basalt in *Artemisia tripartita* and *Artemisia
tridentata wyomingensis* habitat. Site is dominated by native
plants; *Poa* sp., *Phlox* sp., and *Gymnosteris nudicaulis*.

Minimum Elevation: 4774 feet
Maximum Elevation: 4900 feet
Size: 10 AC?

Ownership Comments: Shoshone District BLM, Monument RA.

Comments:

This occurrence was compiled from the following sources: Harrison 1981 (southern population), Packard and Smithman 1984, small-scale (1981-84?) in the Shoshone District BLM Office, Popovich (Shoshone BLM) survey 1993-1994 (northern population).

Protection Comments: Area possibly used for sheep trailing.

Specimens:

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 009

Survey Site Name: BEAR DEN BUTTE

County: Blaine
Minidoka

USGS quadrangle: BEAR DEN BUTTE
BEAR PARK WEST

Latitude: 43 12 33 N Longitude: 113 30 02 W

TOWNRANGE:	SECTION:	TRSNOTE:
002S025E	31	E2SE4
002S025E	32	
002S025E	33	W2W2SW4
003S025E	04	W2W2NW4, S2S2SW4
003S025E	05	
003S025E	06	E2E2
003S025E	08	N2
003S025E	09	N2NW4

Location:
Basin between Bear Den Butte and Lava Butte. About 42 miles SE of
Carey.

Survey Date: 1994-05-11
Last Observed: 1994-05-11
First Observed: 1984

EO RANK: A
EO RANK Comments:

Population Data:
1994: Many thousands of plants of all age classes. Very vigorous
population in high quality habitat. Area surveyed by Bob Moseley,
Idaho CDC.

Habitat Description:
High quality *Artemisia tridentata wyomingensis*/*Stipa comata*
community on deep sandy loess over basalt. Considerable bare
ground. With *Gymnosteris nudicaulis*, *Leptodactylon pungens*,
Bromus tectorum, *Machaeranthera canescens*, *Phlox aculeata*, and
Chrysothamnus nauseosus.

Minimum Elevation: 4840 feet
Maximum Elevation: 5000 feet
Size: 200 + AC

Ownership Comments:
Shoshone District BLM, Monument R.A.

Comments:
The precise population boundaries are unknown. This occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale (1981-84?) in the Shoshone District BLM Office, 7.5' quad map (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Protection Comments:
This is in excellent condition - high quality habitat, extensive population and little (no) grazing pressure in this finger of Laidlaw Park. This site is an excellent candidate for protection as an ACEC.

Specimens:
Lynda C. Smithman LS-1425 (CIC).

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 010

Survey Site Name: ASPITARTE LAKE

County: Minidoka

USGS quadrangle: BEAR DEN BUTTE

Latitude: 43 09 10 N Longitude: 113 35 53 W

TOWNRANGE:	SECTION:	TRSNOTE:
003S024E	10	S2SE4
003S024E	15	
003S024E	16	SE4SE4SE4, SW4SW4SW4
003S024E	17	NE4NE4, N2SE4NE4, S2SE4SE4
003S024E	20	NE4
003S024E	21	N2, NE4SE4
003S024E	22	W2
003S024E	27	N2

Location:

Between Williams Lake and Lava Point, centered on Aspitarte Lake, Laidlaw Park. About 30 mi SE of Carey.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1981-06-10

EORANK: A

EORANK Comments:

Population Data:

1994: Extensive population with good age class distribution, consisting of many thousands of plants. Area surveyed by Bob Moseley, Idaho CDC. 1994: Northwestern population found to be slightly more extensive. Observation by S. Popovich. Population assessed as fair, and no change from last visit.

Habitat Description:

Mostly *Artemisia tridentata wyomingensis*/*Stipa comata* community in rolling terrain. Deep sandy loam over basalt. Associates include *Poa* sp., *Phlox* sp., and *Gymnosteris multicaulis*. Most of the population was not burned by a recent fire, although the southern part of the population is within the area burned by the 1992 Great Rift Fire.

Minimum Elevation: 4580 feet
Maximum Elevation: 4685 feet
Size: 200+ AC

Ownership Comments:
Shoshone District BLM, Monument RA, and state land.

Comments:
The precise population boundaries are unknown. This occurrence was compiled from the following sources: Harrison 1981, Packard and Smithman 1984, small-scale (1981-84?) in the Shoshone District BLM Office, Popovich (Shoshone BLM) survey 1993-1994, and Moseley 1994 survey.

Protection Comments:
The population is locally heavily grazed by cattle due to its proximity to lakes and a well. This site was recommended for protection by Packard and Smithman 1984.

Specimens:
Lynda C. Smithman LS-1418 (CIC).
B. F. and L. C. Harrison 13,428 (Shoshone BLM).

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 011

Survey Site Name: BEAR DEN LAKE

County: Minidoka

USGS quadrangle: BEAR PARK WEST

Latitude: 43 09 56 N Longitude: 113 28 26 W

TOWNRANGE: SECTION: TRSNOTE:
003S025E 16 SE4NE4

Location:
About 41 mi SE of Carey. E of Bear Den Lake.

Survey Date: 1994-05-11
Last Observed: 1984-06-14
First Observed: 1984-06-14

EORANK: H
EORANK Comments: Historical

Population Data:
1984: no data. 1994: Site was not visited by Bob Moseley, Idaho
CDC, during his 1994 survey.

Habitat Description:
Sagebrush-grass zone.

Minimum Elevation: 4840 feet

Size:

Ownership Comments:
State land.

Comments:
This occurrence was compiled from the following sources: Packard
and Smithman 1984, small-scale map (1981-84?) in the Shoshone
District BLM Office, and 7.5' quad (1981-84?) in the Shoshone
District BLM Office.

Specimens:
Lynda C. Smithman LS-1424 (CIC) -- collection label reads
"Laidlaw Park area, Shoshone District BLM."

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 012

Survey Site Name: BEAR DEN LAKE SOUTHEAST

County: Minidoka

USGS quadrangle: BEAR PARK WEST

Latitude: 43 08 36 N Longitude: 113 28 17 W

TOWNRANGE: SECTION: TRSNOTE:
003S025E 22 SW4SW4

Location:

About 44 mi SE of Carey, SE of Bear Den Butte.

Survey Date: 1994-05-11

Last Observed: 1984-06-14

First Observed: 1984-06-14

EORANK: H

EORANK Comments:

Population Data:

1984: no data. 1994: This site was not visited by Bob Moseley,
Idaho CDC, during his 1994 survey.

Habitat Description:

Sagebrush-grass zone.

Minimum Elevation: 4740 feet

Size:

Ownership Comments:

Shoshone District BLM, Monument RA.

Comments:

This occurrence was compiled from the following sources: Harrison
1981 (southern population), Packard and Smithman 1984,
small-scale (1981-84?) in the Shoshone District BLM Office, and
7.5' quad (1981-84?) in the Shoshone District BLM Office.

Specimens:

Lynda C. Smithman LS-1423 (CIC) - collection label reads "Laidlaw
Park area, Shoshone BLM District."

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 013

Survey Site Name: LOWER THUMB RESERVOIR

County: Minidoka

USGS quadrangle: BOTTLENECK LAKE
LAIDLAW LAKE
BEAR DEN BUTTE

Latitude: 43 05 55 N Longitude: 113 32 32 W

TOWNRANGE:	SECTION:	TRSNOTE:
003S024E	31	S2, S2N2
003S024E	32	S2
003S024E	35	E2, SW4
003S024E	36	W2
003S025E	32	SW4
004S024E	01	W2, SW4SE4
004S024E	02	E2, NW4
004S024E	03	N2S2, S2N2
004S024E	04	NW4, S2NE4, N2SE4
004S024E	05	N2NE4
004S025E	06	unsurveyed

Location:

Vicinity of Lower Thumb Reservoir and extending east for ca 3 miles, north for ca 2 miles, and west for ca 5 miles.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1984-06-14

EORANK: A

EORANK Comments:

Population Data:

1994: Tens of thousands of individuals scattered along roughly 10 miles of road. Plants are moderately dense in some areas. Age class distribution is good. Area surveyed by Steve Popovich, Shoshone District BLM, in 1993 and 1994, and by Bob Moseley, Idaho CDC, in 1994. 1994 (Summer): Population assessed as very good to excellent, with a significant increase in recruitment. Cursory visit by S. Popovich, and David Pyke, NBS.

Habitat Description:

Burned (1992) and unburned *Artemisia tridentata wyomingensis*/*Stipa comata* communities with sandy soils. Gently rolling hills. Associates include *Chrysothamnus viscidiflorus*, *Oryzopsis hymenoides*, *Poa secunda*, and *Phlox aculeata*.

Minimum Elevation: 4460 feet

Maximum Elevation: 4640 feet

Size: 500+ AC

Ownership Comments:

Shoshone District BLM, Monument RA, and state land.

Comments:

The boundaries of the population are not precisely known. This occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, 7.5' quad (1981-84?) in the Shoshone District BLM Office, Popovich (Shoshone BLM) surveys 1993-1994, and Moseley 1994 survey.

Protection Comments:

Some areas have dense concentrations of *A. oniciformis* and are worthy of review for special management. Part of the area is grazed by cattle due to proximity of water. Possible threats to population could be decadence of sagebrush stands, and fire rehab reseeding projects in the vicinity.

Management Comments:

Monitoring plots established in T3S R24E Section 35.

Specimens:

Lynda C. Smithman LS-1428 (CIC).

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 014

Survey Site Name: MUD FLAT LAKE NORTHWEST

County: Lincoln

USGS quadrangle: HALFWAY LAKE

Latitude: 43 06 31 N Longitude: 113 46 57 W

TOWNRANGE: SECTION: TRSNOTE:
004S022E 01 W2NE4, E2NW4, N2NW4SE4

Location: About 1 air mile NW of Mud Flat Lake, roughly 26 miles SE of Carey.

Last Observed: 1994
First Observed: 1993

EORANK: A

Population Data:
1993-1994: Area thoroughly surveyed by Steve Popovich, Shoshone District BLM. Ca 1000 plants observed. 1994: Population assessed as very good and a significant increase in recruitment from previous visit. Observation by S. Popovich.

Habitat Description:
In sandier areas and bowl. *Artemisia tridentata wyomingensis* site that was burned in 1992. Site is dominated by native plant species, especially rhizomatous wheatgrass. *Astragalus oniciformis* was not found in nearby areas. Associated spp. include *Stipa comata*, *Oryzopsis hymenoides*, and *Gymnosteris nudicaulis*.

Minimum Elevation: 4620 feet Maximum Elevation: 4640 feet
Size: 40 AC

Ownership Comments: Shoshone District BLM, Monument RA.

Protection Comments: Fire rehab reseeding near site may limit expansion of the population.

Management Comments: Monitoring plots established in the NW4NE4 of Sec 1.

Specimens: Steve Popovich (# and herbarium not yet assigned).

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 015

Survey Site Name: SQUAW BUTTE

County: Minidoka
Lincoln

USGS quadrangle: LAIDLAW LAKE
LAIDLAW BUTTE

Latitude: 43 06 40 N Longitude: 113 39 43 W

TOWNRANGE:	SECTION:	TRSNOTE:
003S023E	20	SW4SE4,SW4NW4SE4
003S023E	21	S2
003S023E	22	S2
003S023E	26	W2, SW4SE4
003S023E	27	N2
003S023E	35	E2, NE4NW4
003S023E	36	S2, SW4NW4
004S023E	01	N2, N2SW4
004S023E	02	S2
004S023E	08	E2

Location:

About 31 mi SE of Carey. Squaw Butte vicinity.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1984

EORANK: A

EORANK Comments: Ranking done in the summer of 1994.

Population Data:

1994: Tens of thousands of plants with good age class distribution. Lower density in areas that were drilled and seeded. Area surveyed by Bob Moseley, Idaho CDC. 1994 (Summer): Population observed extending west and north of previously reported population boundaries. Vigor assessed as excellent on this cursory visit by Steve Popovich, Shoshone District BLM.

Habitat Description:

Area burned in 1992 and was drilled and seeded in 1992. Prominent species include *Poa secunda*, *Lomatium foeniculaceum*, and *Machaeranthera canescens*. Rolling sandy loam over basalt.

Minimum Elevation: 4480 feet
Maximum Elevation: 4640 feet
Size: 500+ AC

Ownership Comments:

Shoshone District BLM, Monument RA, and state land.

Comments:

The precise boundaries of the population are unknown. This occurrence was compiled from the following sources: Packard and Smithman 1984, 7.5' quad in the Shoshone District BLM Office, Popovich (Shoshone BLM) 1993-94 survey, and Moseley 1994 survey.

Management Comments:

Monitoring plots established T3SR23E Section 21.

Specimens:

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 016

Survey Site Name: CRATER BUTTE

County: Lincoln

USGS quadrangle: DIETRICH

Latitude: 42 57 12 N Longitude: 114 17 41 W

TOWNRANGE: SECTION:

005S018E 26

005S018E 27

TRSNOTE:

W2SW4

WSE4, E2SW4, SE4SE4SE4

Location:

West slope of Crater Butte, between the crater and US Route 93,
ca 5.5 miles ENE of Shoshone.

Survey Date: 1994-06-16

Last Observed: 1994-06-16

First Observed: 1980'S

EORANK: C

EORANK Comments:

Population Data:

1994: <100 genets in the NE population; 2000+ genets in the western population ; the southwestern population was not visited in 1994. Plants were 10% in flower and 90% in fruit. Sites north of the fenceline with lower grazing intensities have much higher population densities. Population observed on a cursory visit by Bob Moseley, Idaho CDC. The limits of the western population are unknown.

Habitat Description:

Artemisia tridentata wyomingensis/*Agropyron smithii* habitat with *Chaenactis douglasii*, *Chrysothamnus viscidiflorus*, *Machaeranthera canescens*, and *Phacelia heterophylla*. Gentle slope to the west with deep sandy loam soil. Area north of the fenceline that parallels the road is not heavily grazed and is in better ecological condition. Areas south of the fenceline are in poor condition. Overall site quality is fair.

Minimum Elevation: 4110 feet

Maximum Elevation: 4260 feet

Size: 40 AC

Ownership Comments:
Shoshone District BLM, Monument RA.

Comments:
This occurrence was compiled from the following sources: a small-scale map (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Protection Comments:
The south side of the fence is heavily grazed.

Specimens:

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 017

Survey Site Name: LITTLE WOOD RIVER

County: Lincoln

USGS quadrangle: PAGARI
BLACK RIDGE CRATER
RICHFIELD
TIKURA

Latitude: 43 05 10 N Longitude: 114 05 15 W

TOWNRANGE: SECTION:

003S020E 01,11-14,22,23,26,27,33-36

003S021E 04,06,07,18,31

004S019E 25,36

004S020E 01-04,08-10,15-17,19-21,29-31

Location: Along the Little Wood River upstream from Richfield for ca 12 miles to near the confluence with Silver Creek.

Survey Date: 1994-05-10

Last Observed: 1994

First Observed: 1984-06-26

EORANK: A

EORANK Comments: Ranking dates from spring and fall of 1994.

Population Data:

1994: Many 10,000's of plants. Density depends on community, generally lower in crested wheatgrass seedings. Some areas with high density. Good age class distribution. Area surveyed in 1993 and 1994 by Steve Popovich, Shoshone District BLM, and by Bob Moseley, Idaho CDC, in 1994. 1994 (Spring): Additional subpopulation east of first population reported. 400 genets observed on a cursory visit by Caryl Elzinga, Alderspring Ecological Consulting. Subpopulation vigor assessed as fair. 1994 (Fall): Additional subpopulations observed by S. Popovich and David Pyke, NBS. Vigor assessed as good to excellent.

Habitat Description:

Mostly *Artemisia tridentata wyomingensis*/*Stipa comata* communities but also some *Agropyron cristatum* seedings. Sandy loam soil over basalt. Rolling terrain - flat to gentle slopes. Generally considerable bare ground in communities.

Minimum Elevation: 4260 feet
Maximum Elevation: 4650 feet
Size: 500+ AC

Land Owner/Manager:
SHOSHONE DISTRICT BLM, MONUMENT RA
BEAR TRACK WILLIAMS RECREATION AREA
STATE LAND WITHIN LINCOLN COUNTY
PRIVATE LAND

Comments:

The exact boundaries of the occurrence are unknown, but it is more or less continuous in the area. The western boundary coincides with US Route 93, west of which are clayey soils unsuitable for *A. oniciformis*. The eastern boundary is unknown. This occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, Popovich 1993-1994 surveys, and Moseley 1994 survey.

Protection Comments:

Some sites are fairly high in quality, but most sites are grazed by cattle and/or sheep and are very weedy. There are currently seeded areas, with future seedings possible. Many roads traverse the occurrence as well as an abandoned railroad right-of-way.

Management Comments:

Monitoring plots established T4SR20E Section 8 (at mile marker 186 on Highway 26), and at several locations along road east of Pagari Bridge.

Specimens:

Lynda C. Smithman LS-1494 (CIC).
Lynda C. Smithman LS-1495 (CIC).
Lynda C. Smithman LS-1496 (CIC).
Lynda C. Smithman LS-1497 (CIC).
Steve Popovich (# and herbarium not yet assigned).
Caryl Elzinga 6371 (herbarium not yet assigned).

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 018

Survey Site Name: THE BLOW OUT SOUTHWEST

County: Lincoln

USGS quadrangle: WAGON BUTTE

Latitude: 43 07 31 N Longitude: 113 47 02 W

TOWNRANGE: SECTION: TRSNOTE:
003S022E 36 NW4NE4

Location:

About 1.5 air miles SW of The Blow Out, 2.8 miles S of Cut Off Reservoir, and roughly 25 air miles SE of Carey.

Survey Date:

Last Observed: 1994-05-16

First Observed: 1993-1994

EORANK: C

EORANK Comments:

Population Data:

1993-1994: Area surveyed by Steve Popovich, Shoshone District BLM. About 100 plants observed. 1994: 1994: 100 genets observed by S. Popovich on a thorough survey. Population assessed as fair.

Habitat Description:

In sandy swale that burned in 1992. With *Agropyron dasytachyum*, *Carex* sp., and *Phlox* sp. No *A. oniciformis* was found in nearby areas.

Minimum Elevation: 4640 feet

Size: 1 AC

Ownership Comments:

State land.

Protection Comments:

Possible threats may be repeated fire and grazing.

Specimens:

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 019

Survey Site Name: MONUMENT BUTTE

County: Lincoln

USGS quadrangle: PAGARI WELL

Latitude: 43 09 35 N Longitude: 113 56 20 W

TOWNRANGE:	SECTION:	TRSNOTE:
003S021E	15	E2SE4
003S021E	16	E2NE4
003S021E	17	W2SE4,S2SE4SW4
003S021E	20	W2NE4,E2NW4

Location:

About 12 miles SW of Carey; E of US Route 93 ca 15 miles; within a 1 mile radius NE, N, and NW of Monument Butte.

Survey Date: 1994-05-10

Last Observed: 1994-05-19

First Observed: 1984

EORANK: C

EORANK Comments:

Population Data:

1994: Hundreds of plants in a low density population. Area surveyed by Bob Moseley, Idaho CDC. 1994: Ca 100 additional plants observed in 2 subpopulations west of the originally reported populations. Plant vigor assessed as good on this cursory visit by Caryl Elzinga, Alderspring Ecological Consulting.

Habitat Description:

Area has been converted to *Agropyron spicatum* or is heavily grazed due to the presence of a water development.

Minimum Elevation: 4600 feet

Maximum Elevation: 4660 feet

Size: 40? AC

Ownership Comments:

Shoshone District BLM, Monument RA, and state land.

Comments:

The exact population boundaries are unknown. This occurrence was compiled from the following sources: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, 7.5' quad (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

The area has been overseeded with *Agropyron* spp.; grazing and weed invasion also exist.

Specimens:

Caryl Elzinga 6370 (herbarium unknown).

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 020

Survey Site Name: BACON LAKE

County: Lincoln

USGS quadrangle: WAGON BUTTE

Latitude: 43 09 43 N Longitude: 113 46 47 W

TOWNRANGE:	SECTION:	TRSNOTE:
003S022E	13	SE4, SW4NE4
003S023E	18	E2SW4, W2SE4

Location:

About 22 miles S of Carey on the road S of Wagon Butte in the vicinity of Bacon Lake.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1984

EORANK: C

EORANK Comments:

Population Data:

1994: Hundreds of plants in a population of moderate density. Good age class distribution. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Unburned *Artemisia tridentata wyomingensis*/*Stipa comata* community in fair condition. Lots of *Bromus tectorum*.

Minimum Elevation: 4700 feet Maximum Elevation: 4720 feet
Size: 20? AC

Ownership Comments:

Shoshone District BLM, Monument RA.

Comments:

The precise population boundaries are unknown. This occurrence was compiled from the following: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, 7.5' quad map (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Protection Comments:

Area is heavily grazed due to the presence of water developments in the area.

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 021

Survey Site Name: THE BLOW OUT WEST

County: Lincoln

USGS quadrangle: WAGON BUTTE

Latitude: 43 08 23 N Longitude: 113 47 18 W

TOWNRANGE: SECTION: TRSNOTE:
003S022E 25 center N2NW4

Location:

Ca 1 air mile WSW of The Blow Out and ca 24 miles SE of Carey.

Survey Date:

Last Observed: 1994-05-16

First Observed: 1993-1994

EORANK: D

EORANK Comments:

Population Data:

1993-1994: Area surveyed by Steve Popovich, Shoshone District BLM. About 100 plants observed. 1994: Area thoroughly surveyed by S. Popovich. Population assessed as fair.

Habitat Description:

Sandy plain that burned in 1992. With *Agoseris* sp., *Poa* sp., *Sisymbrium altissimum*, *Artemisia tripartita*, and *Chrysothamnus* sp. No *Astragalus oniciformis* was found in nearby areas.

Minimum Elevation: 4670 feet

Size: 3 AC

Ownership Comments:

Shoshone District BLM, Monument RA.

Protection Comments:

Possible threats may be repeated fire and grazing, and invasion of annuals.

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 022

Survey Site Name: THE BLOW OUT

County: Lincoln

USGS quadrangle: WAGON BUTTE

Latitude: 43 08 55 N Longitude: 113 46 49 W

TOWNRANGE: SECTION:
003S022E 24

TRSNOTE:
NW4SE4, SW4NE4

Location:

Just W of The Blow Out and roughly 23 air miles SE of Carey.

Survey Date:

Last Observed: 1994-05-16

First Observed: 1993-1994

EORANK: D

EORANK Comments:

Population Data:

1993-1994: Area surveyed by Steve Popovich, Shoshone District BLM. About 30 plants observed. 1994: 30 genets; population assessed as having very little recruitment, but good, large, mature individuals. area thoroughly surveyed by S. Popovich.

Habitat Description:

A sandy site with unburned, decadent *Artemisia tridentata wyomingensis* and *Artemisia tripartita* with *Poa* sp. *Astragalus oniciformis* is mostly confined to open interstitial areas. No plants were found in nearby burned area with finer textured soil.

Minimum Elevation: 4690 feet

Size: 3 AC

Ownership Comments: Shoshone District BLM. Monument RA.

Protection Comments:

Possible threats may be repeated fire and grazing, and stagnation of the sage stand.

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 023

Survey Site Name: STUDENT LAKE

County: Blaine

USGS quadrangle: BEAR DEN BUTTE

Latitude: 43 13 30 N Longitude: 113 31 40 W

TOWNRANGE:	SECTION:	TRSNOTE:
002S024E	25	SE4, SE4SW4
002S024E	36	NW4
002S025E	18	SE4SE4
002S025E	19	NE4NE4
002S025E	30	NW4

Location:

Along road N and S of Student Lake in Laidlaw Park.

Survey Date: 1994-05-11

Last Observed: 1994-05-11

First Observed: 1980'S

EORANK: A

EORANK Comments:

Population Data:

1994: Thousands of flowering genets. Age class distribution is good. Population are locally dense. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Burned and unburned *Artemisia tridentata wyomingensis/Stipa comata* stands with mostly native species, although some *Sisymbrium altissimum* and *Bromus tectorum* are present. Sandy loam soil over basalt with considerable bare ground. Flat to rolling slopes. Good overall site quality.

Minimum Elevation: 4860 feet

Maximum Elevation: 5000 feet

Size: 40 + AC

Ownership Comments:

Shoshone District BLM, Monument RA, and state land.

Comments:

The precise boundaries of the population are unknown. This occurrence was compiled from the following sources: small-scale map (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 024

Survey Site Name: MULE BUTTE

County: Blaine

USGS quadrangle: MULE BUTTE

Latitude: 43 03 58 N Longitude: 113 21 38 W

TOWNRANGE:	SECTION:	TRSNOTE:
004S026E	13	E2SE4SE4; UNSURVEYED
004S026E	24	E2NE4NE4; UNSURVEYED
004S027E	18	W2SW4
004S027E	19	NW4

Location:

Mule Butte; due W of the bowl of Mule Butte to 0.5 miles S of the bowl of Mule Butte.

Survey Date: 1994-06-22

Last Observed: 1994-06-22

First Observed: 1994-06-22

EORANK: A

EORANK Comments:

Population Data:

1994: 1000-10,000 genets estimated; 5% vegetative, 5% flowering, 85% in fruit, and 5% dormant. Age class structure was 5% seedlings, 20% immature, 70% mature, and 5% senescent. Population vigor assessed as very healthy, with good fruit production. Cursory visit by Steve Popovich and Sharon Weiss, Shoshone District BLM.

Habitat Description:

Occurs in gentle rolling *Artemisia tridentata wyomingensis* basalts of the Snake River Plains, in sandy soil areas of grassy openings dominated by *Stipa comata*, rhizomatous *Agropyron* sp., with *Oryzopsis hymenoides*, *Poa* spp., *Stipa thurberiana*, *Phacelia* sp., *Opuntia* sp., *Eriogonum* sp., *Bromus tectorum*, *Alyssum desertorum*, *Machaeranthera canescens*, *Astragalus purshii*, and *A. lentiginosus*.

Minimum Elevation: 4810 feet

Maximum Elevation: 4980 feet

Size: 40 AC

Ownership Comments:

Idaho Falls District BLM, Big Butte RA.

Protection comments:

Land is currently used for light sheep trailing. Possible threats may be potential frequent fires.

Specimens: S. Popovich (# and herbarium not yet assigned).

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 025

Survey Site Name: WAY EAST OF RICHFIELD

County: Lincoln

USGS quadrangle: PAGARI

Latitude: 43 02 03 N Longitude: 114 01 23 W

TOWNRANGE: SECTION:
004S020E 36

Location:

About 20 mi S of Carey, turn E off US Route 26/93 and head SE
nearly 6 mi to jeep trail SW which heads into Sec. 36

Survey Date: 1994-05-11

Last Observed: 1984

First Observed: 1984

EORANK: H

EORANK Comments:

Population Data:

1984: No data. 1994: Bob Moseley, Idaho CDC, was unable to
relocate the population. There is little habitat in this section.

Habitat Description:

Minimum Elevation: 4325 feet

Size:

Ownership Comments:

State land.

Comments:

Packard and Smithman (1984) list *A. oniciformis* as
occurring in this section.

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 026

Survey Site Name: BLACK RIDGE CRATER NORTH

County: Lincoln

USGS quadrangle: BLACK RIDGE CRATER

Latitude: 43 04 47 N Longitude: 113 56 25 W

TOWNRANGE:	SECTION:	TRSNOTE:
004S021E	15	NE4, SW4NW4
004S021E	16	SE4NE4

Location:

About 17 mi S of Carey turn SE from US Route 26/93 (over Pagari Bridge) and head SE approximately 7 mi.

Survey Date: 1994-05-10

Last Observed: 1994-05-10

First Observed: 1984

EORANK: B

Population Data: 1994: Dense population of thousands of individuals with good age class distribution. The western population was not visited in 1994. Surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Depression in lava with deep sands. Surrounding lava plains have little sandy loess. Area burned in 1992. *Artemisia tridentata wyomingensis*/*Stipa comata* community with *Oryzopsis hymenoides*, *Phacelia heterophylla*, and *Chrysothamnus viscidiflorus*. Considerable bare ground.

Minimum Elevation: 4500 feet Maximum Elevation: 4520 feet
Size: 10 AC

Ownership Comments: Shoshone District BLM, Monument RA, and state land.

Comments: The exact population boundaries are unknown. This occurrence was compiled from the following: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Specimens:

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 027

Survey Site Name: BLACK RIDGE CRATER

County: Lincoln

USGS quadrangle: BLACK RIDGE CRATER

Latitude: 43 01 57 N Longitude: 113 59 47 W

TOWNRANGE:	SECTION:	TRSNOTE:
004S021E	31	NE4

Location:

About 20 mi S of Carey turn E from US Route 26/93 and head SE over 7 mi. NW of Black Ridge Crater.

Survey Date: 1994-05-12

Last Observed: 1984

First Observed: 1984

EORANK: H

EORANK Comments: Historical

Population Data:

1994: Bob Moseley, Idaho CDC, did not visit this site during the 1994 survey for *A. oniciformis*.

Habitat Description:

Minimum Elevation: 4440 feet

Size:

Ownership Comments:

Shoshone District BLM, Monument RA.

Comments:

Packard and Smithman (1984) list the population as being in section 31. No other data is given. This occurrence was compiled from the following: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, and 7.5' quad (1981-84?) in the Shoshone District BLM Office.

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 028

Survey Site Name: TUNUPA

County: Lincoln

USGS quadrangle: TUNUPA

Latitude: 42 56 45 N Longitude: 114 32 38 W

TOWNRANGE: SECTION:
005S016E 34

Location:

About 5.8 mi W of Shoshone (10 mi E Gooding) turn NW from US
Route 20/26 and proceed W for about 1 mile.

Survey Date: 1994-06-16

Last Observed: 1984

First Observed: 1984

EORANK: H

EORANK Comments: Historical

Population Data:

1994: Bob Moseley, Idaho CDC, was unable to relocate the
population.

Habitat Description:

Minimum Elevation: 3770 feet

Size:

Ownership Comments:

Shoshone District BLM, Bennett Hills RA or private land.

Comments:

Occurrence information compiled from the following sources:
Packard and Smithman 1984, 7.5' quad (1981-84?) in Shoshone BLM
Office.

Protection Comments:

Specimens:

ASTRAGALUS ONICIFORMIS
PICABO MILKVETCH
Occurrence Number: 029

Survey Site Name: MULE BUTTE NORTHEAST

County: Blaine

USGS quadrangle: MULE BUTTE

Latitude: 43 05 17 N Longitude: 113 19 22 W

TOWNRANGE: SECTION: TRSNOTE:
004S027E 08 NE4NE4SE4

Location:

2.1 airmiles NE of Mule Butte, 0.3 miles west of 3-way intersection, along both sides of the jeep trail.

Survey Date: 1994-06-23

Last Observed: 1994-06-23

First Observed: 1994-06-23

EORANK: D

Population Data:

1994: Less than 50 genets estimated, 2% in flower and 98% in fruit. Age class structure was 5% immature, 50% mature, and 45% senescent. Population vigor assessed as poor on this cursory visit by Steve Popovich and Sharon Weiss, Shoshone District BLM.

Habitat Description:

5% slope, SE aspect. In sandy soils over basalt rolling hills of Snake River Plains. Found growing in open areas dominated by rhizomatous wheatgrass (may indicate repeated fire or grazing). Associated species include *Poa* sp., *Phacelia* sp., *Artemisia tridentata wyomingensis*, *Chrysothamnus* sp., *Opuntia* sp., *Stipa comata*, *Bromus tectorum*, *Machaeranthera canescens*, *Alyssum desertorum* and *Phlox* sp. Site is in the early seral stage.

Minimum Elevation: 4890 feet Size: 2 AC

Ownership Comments: Idaho Falls District BLM. Big Butte RA.

Protection Comments: There is evidence of trailing by sheep; also a recent burn with a high conversion to annual range.

Specimens:

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 030

Survey Site Name: BROWN BUTTE

County: Lincoln

USGS quadrangle: DIETRICH BUTTE
RICHFIELD

Latitude: 42 59 18 N Longitude: 114 11 28 W

TOWNRANGE:	SECTION:	TRSNOTE:
005S019E	10	SE4
005S019E	11	N2
005S019E	15	W2
005S019E	16	E2NE4, NE4SE4
005S019E	20	NE4SE4, SE4NE4
005S019E	21	W2SW4, SW4NW4
005S019E	22	NW4

Location:

About 6.5 mi S of Richfield turn off US Route 20/26/Alt 93 to the E and proceed past Brown Butte. Several subpopulations observed N of Dietrich Butte on both sides of the Dietrich Main Canal for ca 2.5 miles. E of Brown Butte.

Survey Date: 1994-05-13

Last Observed: 1994-05-13

First Observed: 1984-06-27

EORANK: C

EORANK Comments:

Population Data:

1994: Thousands of individuals apparently with good age class distribution. SW subpopulation not visited during the 1994 survey. Area surveyed by Bob Moseley, Idaho CDC.

Habitat Description:

Sandy soils; probably *Artemisia tridentata wyomingensis*/*Stipa comata* habitat type degraded by heavy cattle grazing and now containing an unnaturally high ground cover of weeds. Some areas converted to or interplanted with *Agropyron cristatum*. *Bromus tectorum* has higher cover here than other sites to the N and NE.

Minimum Elevation: 4170 feet Maximum Elevation: 4240 feet

Size: 10 AC

Ownership Comments:

Shoshone District BLM, Monument RA.

Comments:

The exact boundaries of the occurrence are unknown. It is probably more continuous than currently mapped. The occurrence was compiled from the following: Packard and Smithman 1984, small-scale map (1981-84?) in the Shoshone District BLM Office, and Moseley 1994 survey.

Protection Comments:

Habitat was destroyed by construction roads and the Dietrich Main Canal. Habitat is degraded by cattle grazing, which is ongoing.

Specimens:

Lynda C. Smithman LS-1498 (CIC).

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 031

Survey Site Name: SAND DUNES

County: Lincoln

USGS quadrangle: DIETRICH BUTTE

Latitude: 42 53 17 N Longitude: 114 08 30 W

TOWNRANGE: SECTION: TRSNOTE:
006S019E 24 SW4NE4

Location:
Off State Route 24, sand dune area 7.5 mi E of Dietrich (19 mi W
of Kimama).

Survey Date: 1994-06-16
Last Observed: 1984-06-12
First Observed: 1984-06-12

EO RANK: H
EO RANK Comments: Historical

Population Data:
1984: no data. 1994: Bob Moseley, Idaho CDC, was unable to
relocate the population. Most of the area supports *Psoralea*
lanceolata, indicating a habitat too sandy for *A. oniciformis*.

Habitat Description:
Sandy loam at edge of dune area and on surrounding plains.
Artemisia tridentata; stabilized habitat.

Minimum Elevation: 4170 feet

Size:

Ownership Comments:
Shoshone District BLM, Monument RA.

Comments:
Occurrence compiled from the following sources: Packard and
Smithman 1984, small scale map (1981-84?) in Shoshone BLM Office,
and Moseley 1994 summary.

Specimens: Lynda C. Smithman LS-1402 (CIC).

· *ASTRAGALUS ONICIFORMIS*

PICABO MILKVETCH

Occurrence Number: 032

Survey Site Name: GREAT RIFT #1

County: Blaine

USGS quadrangle: MULE BUTTE

Latitude: 43 06 22 N Longitude: 113 19 04 W

TOWNRANGE: SECTION: TRSNOTE:
004S027E 4 NE4SW4NW4

Location:

3 air miles NE of Mule Butte; just south of a faint intersection of jeep trails.

Survey Date: 1994-06-23

Last Observed: 1994-06-23

First Observed: 1994-06-23

EORANK: B

Population Data:

1994: Ca 300 genets observed; 10% vegetative, 10% flower, 75% fruit, and 5% dormant. Population age class structure was 10% seedlings, 60% immature, 20% mature, and 10% senescent. Vigor assessed as good with a lot of juveniles present. Cursory visit by Steve Popovich and Sharon Weiss, Shoshone District BLM.

Habitat Description:

In silty, sandy, or clay soils over basalts of Snake River Plains; occurs in bowl or depression areas, along roadsides, or in more open areas with *Bromus tectorum*, *Alyssum desertorum*, *Artemisia tridentata wyomingensis*, *Chrysothamnus* sp., *Machaeranthera canescens*, *Phlox* sp., *Poa secunda*, *Stipa comata*, and *Agropyron* sp.

Minimum Elevation: 4810 feet Size: 3 AC

Ownership Comments: Idaho Falls District BLM, Big Butte RA.

Protection Comments: Currently the area is used for light sheep trailing; possible threats could be frequent fires.

Specimens:

·*ASTRAGALUS ONICIFORMIS*
PICABO MILKVETCH
Occurrence Number: 033

Survey Site Name: GREAT RIFT #2

County: Blaine

USGS quadrangle: MULE BUTTE
BEAR PARK EAST

Latitude: 43 07 17 N Longitude: 113 20 05 W

TOWNRANGE:	SECTION:	TRSNOTE:
003S027E	29	S2SE4SW4
003S027E	32	E2NW4

Location:

3.4 airmiles NNE of Mule Butte; on both sides of jeep trail partly surrounding a conspicuous playa.

Survey Date: 1994-06-23

Last Observed: 1994-06-23

First Observed: 1994-06-23

EORANK: A

EORANK Comments: Ranking ranges from excellent to good.

Population Data:

1994: 1100 genets estimated in two populations; of all age classes and in vegetative, flowering, and fruiting stages. Population vigor assessed as fair to very healthy with lots of juveniles and fruit present. Mature plants to 1.5 feet in diameter. Observation on a cursory visit by Steve Popovich and Sharon Weiss, Shoshone BLM.

Habitat Description:

Flat or gentle bowl areas of silt, clay, or sandy soils. Occurs in grassy openings of rolling *Artemisia tridentata wyomingensis* on basalts of Snake River Plains. Associated spp. include *Stipa comata*, *Agropyron* sp., *Poa* sp., *Machaeranthera canescens*, *Alyssum desertorum*, and *Phlox* sp.

Minimum Elevation: 4780 feet

Size: 11 AC

Ownership Comments:

Idaho Falls District BLM, Big Butte RA.

Protection Comments:

Potential frequent fires could lead to strong invasion of weedy annuals, sheep could currently be using the playa as a watering station.

Specimens:

S. Popovich (# and herbarium not yet assigned).

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 034

Survey Site Name: TUNUPA SOUTH

County: Lincoln

USGS quadrangle: TUNUPA

Latitude: 42 56 41 N Longitude: 114 34 44 W

TOWNRANGE: SECTION: TRSNOTE:
005S016E 32 NE4SE4

Location:

Approx 6 miles E of Gooding and 25 mi NW of Twin Falls; S of Tunupa and the Little Wood River.

Survey Date: 1994-06-16

Last Observed: 1989-05-12

First Observed: 1989

EORANK: D

EORANK Comments:

Population Data:

1989: 50-100 individuals in flower and fruit; poor overall site quality. Observation by Ann DeBolt, Boise District BLM. 1994: Bob Moseley, Idaho CDC, was unable to relocate the population.

Habitat Description:

Sandy soils in basin *Artemisia tridentata* habitat where *Bromus tectorum* has become prevalent.

Minimum Elevation: 3700 feet

Size: 2 AC

Ownership Comments:

Shoshone District BLM, Bennett Hills RA.

Comments: Observation report gives elevation as 3100 feet, which is too low for the location described.

Protection Comments: Currently grazed; population is threatened by range fires.

Specimens: A. DeBolt 1111 (herbarium unknown)

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 035

Survey Site Name: CRATER BUTTE SOUTHWEST

County: Lincoln

USGS quadrangle: DIETRICH

Latitude: 42 56 23 N Longitude: 114 19 43 W

TOWNRANGE: SECTION: TRSNOTE:
005S018E 33 S2SW4

Location:

Just S of the Little Wood River and on either side of US Route 20/26, ca 2.5 air miles NW of Dietrich.

Survey Date: 1994-06-16

Last Observed: 1994-06-16

First Observed: 1992-07-02

EORANK: D

EORANK Comments:

Population Data:

1992: ca 200 plants, all past fruit, occurring in scattered clusters of 4-10 plants each. Observation by Steve Popovich, Shoshone District BLM. 1994: Area surveyed by Bob Moseley, Idaho CDC. No change in population observed.

Habitat Description:

Sandy soil. Poor condition *Artemisia tridentata* area infested by *Bromus tectorum*. Annual *Eriogonum* sp. present.

Minimum Elevation: 4035 feet

Maximum Elevation: 4070 feet

Size: 100 AC

Ownership Comments:

Shoshone District BLM, Monument RA; population may extend onto adjacent private land.

Protection Comments:

The area is disturbed by grazing.

Specimens:

ASTRAGALUS ONICIFORMIS

PICABO MILKVETCH

Occurrence Number: 036

Survey Site Name: DITCH SPRING

County: Blaine

USGS quadrangle: TIKURA

Latitude: 43 14 24 N Longitude: 114 07 07 W

TOWNRANGE: SECTION: TRSNOTE:
002S020E 19 NE4

Location:

Ca 0.8 and 1 mile WSW of Ditch Spring, just S of the road near the base of the S slopes of the Picabo Hills.

Survey Date: 1994-05-12

Last Observed: 1994-05-12

First Observed: 1992-05-14

EORANK: C

Population Data: 1992: 2 subpopulations, each containing 10 plants, 50% in flower, 50% in fruit on 5/14, 100% in fruit by 6/24. Observation was made on a cursory survey by Steve Popovich, Shoshone District BLM. 1994: Localized to these two knobs on both sides of the road. Several hundred plants at each site with good age class distribution. Observation by Bob Moseley, Idaho CDC.

Habitat Description:

Sandy areas around knobs. This is the only known occurrence that is on the hills, not on volcanic plain. Deep sandy loam. Fair to moderate (20%) slopes to the N, E, and S. Associated with *Lupinus*, *Chaenactis* spp., *Collinsia parviflora*, *Mimulus nanus*, *Purshia tridentata*, *Artemisia tridentata wyomingensis*, and annual *Eriogonum* sp.

Minimum Elevation: 4760 feet Maximum Elevation: 4795 feet

Size: 2 AC

Ownership Comments: Shoshone District BLM, Monument RA.

Comments: The site is worthy of further investigation for other rare plants, especially annuals.

Specimens:



Bureau of Land Management
Idaho State Office
3380 Americana Terrace
Boise, Idaho 83706

BLM/ID/PT-95/013+1150